



# Therapeutic Potential of Methanolic Stem Extract of *Costus afer* on Serum Uric Acid, Tumor Necrosis Factor-alpha ( $TNF-\alpha$ ), and Interleukin-6 ( $IL-6$ ) Levels in Monosodium Urate (MSU)-Induced Gouty Arthritis in Wistar Rats

Kpomah Enyohwo Dennis <sup>a\*</sup> and Owo Gogo James <sup>b</sup>

<sup>a</sup> Department of Biochemistry, Federal University, Otuoke, Bayelsa State, Nigeria.

<sup>b</sup> Department of Chemistry, Faculty of Natural and Applied Sciences, Ignatius Ajuru University of Education, Rumuolumeni, Port Harcourt, Rivers State, Nigeria.

## Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jamps/2024/v26i11724>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124422>

Original Research Article

Received: 03/08/2024

Accepted: 08/10/2024

Published: 15/10/2024

## ABSTRACT

The therapeutic potential of methanolic stem extracts of *Costus afer* was investigated to evaluate their effects on serum uric acid,  $TNF-\alpha$ , and  $IL-6$  levels, which are the key biomarkers associated with inflammatory and metabolic disorders like gouty arthritis. Qualitative and quantitative

\*Corresponding author: E-mail: [kpomahed@fuotuoake.edu.ng](mailto:kpomahed@fuotuoake.edu.ng);

**Cite as:** Dennis, Kpomah Enyohwo, and Owo Gogo James. 2024. "Therapeutic Potential of Methanolic Stem Extract of *Costus Afer* on Serum Uric Acid, Tumor Necrosis Factor-Alpha ( $TNF-\alpha$ ), and Interleukin-6 ( $IL-6$ ) Levels in Monosodium Urate (MSU)-Induced Gouty Arthritis in Wistar Rats". *Journal of Advances in Medical and Pharmaceutical Sciences* 26 (11):45-54. <https://doi.org/10.9734/jamps/2024/v26i11724>.

phytochemical analysis of *C. afer* indicated the presence of flavonoids, alkaloids, phenol, anthraquinone and glycosides, these bioactive compounds have been implicated for numerous medicinal uses and as such the therapeutic potentials of the extract were investigated by administering varying doses (100 mg/kg, 200 mg/kg, and 400 mg/kg) of the extract to a Wistar rat model with induced gouty arthritis and assessing its impact compared to a control group and an untreated MSU group. The results revealed that the extracts significantly reduced ( $p < 0.05$ ) serum uric acid levels in a dose-dependent manner, with the high dose achieving a reduction below the control level. Additionally, the extracts effectively decreased ( $p < 0.05$ ) *TNF- $\alpha$*  and *IL-6* levels, indicating potent anti-inflammatory properties. The high dose exhibited the most substantial reduction in these biomarkers, underscoring the extract's therapeutic potential. These findings support the traditional use of *C. afer* and provide scientific validation for its application in managing hyperuricemia and inflammatory conditions.

**Keywords:** *Costus afer*; serum uric acid; *TNF- $\alpha$* ; *IL-6*; anti-inflammatory; hyperuricemia; therapeutic potential.

## 1. INTRODUCTION

Gouty arthritis is a type of inflammatory arthritis that is fundamentally linked to disturbances in purine metabolism and uric acid levels. It is characterized by the deposition of urate crystals in the joints and soft tissues, which leads to intense inflammatory reactions [1]. Gouty arthritis, characterized by elevated serum uric acid levels (hyperuricemia) and severe joint inflammation, poses significant health challenges [2,3]. The condition is often associated with the accumulation of monosodium urate (MSU) crystals in the joints. The presence of MSU crystals in the joints activates the *NLRP3* inflammasome in macrophages, leading to the release of pro-inflammatory cytokines such as interleukin-6 (*IL-6*) and tumour necrosis factor- $\alpha$  (*TNF- $\alpha$* ) [4]. *TNF- $\alpha$*  and *IL-6* are both cytokines involved in inflammation and immune responses [5,6]. *TNF- $\alpha$*  is a pro-inflammatory cytokine produced by various cell types, including macrophages and *T-cells*, while *IL-6* on the other hand, is produced by various cells, including macrophages, *T-cells*, and fibroblasts [7]. *TNF- $\alpha$*  plays a key role in systemic inflammation and is involved in the regulation of immune cells [6,8]. Elevated levels of *TNF- $\alpha$*  are associated with various inflammatory diseases and conditions [9]. Likewise, *IL-6* levels are often elevated in response to infection, injury, or inflammation [10]. Both *TNF- $\alpha$*  and *IL-6* result in an acute inflammatory response in gouty arthritis, characterized by pain, redness, warmth, and swelling in the affected joints. Chronic cases of gouty arthritis may involve the formation of tophi and persistent joint damage [11]. They are targets for therapeutic intervention in gouty arthritis [12]. Therapeutic agents that inhibit

these cytokines can help reduce inflammation and alleviate symptoms [13].

Over the years the vast treasures inherent in medicinal plants are constantly harnessed to treat and manage diverse health challenges plaguing mankind. These nature's gift to mankind is rich in phytochemicals [14]. The increased patronage of plant medicine arises from the fact that they are perceived as having minimal side effects [15] in comparison to synthetic drugs [16,17] nearly all year round availability [18], and general affordability because of their low cost [19]. *C. afer* has been studied for its potential health benefits, including its antimicrobial [20], anti-inflammatory [21], and antioxidant properties [22]. These attributes make it a subject of interest for investigating its efficacy in treating inflammatory diseases like gouty arthritis. *Costus afer* is indigenous to the people of West Africa, the plant is popularly called ginger lily or bush cane. It is a tall, perennial, and rhizomatous herb. It belongs to the Family of Zingiberaceae presently known as Costaceae it is mostly found in moist or shady forests and river banks [21].

This study aims to evaluate the therapeutic potential of methanolic stem extracts of *C. afer* on inflammatory markers *TNF- $\alpha$* , *IL-6* and serum uric acid levels in Wistar rats with MSU-induced gouty arthritis.

## 2. MATERIALS AND METHODS

### 2.1 Plant Material

Fresh stems of *C. afer* were procured from a local market in Port Harcourt, Rivers State, Nigeria. The plant material was authenticated to confirm its species identity at the herbarium unit

of the Biology Department, Federal University Otuoke, Bayelsa, Nigeria. After authentication, the stems were air-dried in the shade to preserve the integrity of the active compounds. Once dried, the stems were ground into a fine powder using a mechanical grinder. The powdered material was then stored in airtight containers to maintain its quality and prevent moisture absorption until it was needed for extraction.

## 2.2 Extraction Procedure

The cold extraction method was employed for the extraction, a 200 g sample of the powdered stem material was subjected to extraction using methanol (1:5 w/v) for 72 hours at room temperature. The mixture was filtered, and the solvent was evaporated using a rotary evaporator. The resulting extract was stored at 4°C until use.

## 2.3 Phytochemical Analysis

### 2.3.1 Qualitative phytochemical screening

Qualitative phytochemical screening of the methanolic stem extracts of *C. afer* was conducted to identify the presence of various bioactive compounds. The screening included tests for flavonoids, tannins, saponins, alkaloids, phenols, anthraquinones, cardiac glycosides, terpenoids, phlobatannins, and cardenolides. These tests were performed following the standard procedures outlined by Anyasor et al. [22].

### 2.3.2 Quantitative phytochemical analysis

The procedures for the quantitative determination of tannins, alkaloids, flavonoids, saponins, and phenols, were adapted from the methods described by Kpomah et al. [23]. The procedures for the quantitative determination of anthraquinones, cardiac glycosides, terpenoids, phlobatannins, and cardenolides were adapted from the methods described by Wu et al. [24], Ikeda et al. [25] and Malik [26] respectively.

### 2.3.3 Antioxidant activity

The antioxidant activity was estimated using the procedure described by Anyasor et al. [22]. Rapid thin layer chromatography (TLC) was employed by applying a concentrated methanolic solution of the extract onto silica gel plates. These plates were then developed using methanol: ethyl acetate mixture (2:1), allowed to

air dry and sprayed with a 0.2% w/v DPPH solution in methanol. The presence of yellow spots on the plates indicated antioxidant activity. For further evaluation, the DPPH radical scavenging activity of the extracts was measured using the spectrophotometric method outlined by Anyasor et al. [22]. Specifically, 1 mL of a 0.3 mM DPPH methanol solution was mixed with 2.5 mL of the extract or standard solutions (at concentrations of 100, 200, 300, and 400 µg/mL) and incubated at room temperature for 30 minutes. The absorbance of the resulting solution was recorded at 518 nm and used to calculate the percentage antioxidant activity (AA %), using the formula:

$$AA \% = \frac{[100 - ((Abs_{sample} - Abs_{blank}) \times 100)]}{Abs_{control}} \quad (1)$$

Methanol (1.0 mL) combined with the extract solution (2.5 mL) served as the blank, while 1 mL of 0.3 mM DPPH plus methanol (2.5 mL) was the negative control. Ascorbic acid and gallic acid were used as positive controls. The assay was conducted in triplicate for each concentration. The IC<sub>50</sub> values for methanol extract, aqueous extract, ascorbic acid, and gallic acid were calculated.

## 2.4 Animal Model

Twenty five Wistar rats averagely weighing 235.67 ± 6.88 g were used for the study. The rats were acclimatized for one week under standard laboratory conditions prior to the experiment, the animals were housed under controlled conditions with a 12-hour light/dark cycle and provided with standard rodent chow and water *ad-libitum*. The study was approved by the Directorate of Research and Quality Assurance, Federal University Otuoke, Bayelsa State vide a letter referenced DRQA/FUO/0113/11/06/24.

## 2.5 Induction of Gouty Arthritis

Gouty arthritis was induced in rats by intra-articular injection of monosodium urate (MSU) crystals (0.1 mg/rat) into the knee joint under anesthesia (ketamine 60 mg/kg and xylazine 10 mg/kg).

## 2.6 Experimental Design

The rats were divided into five groups (n = 5 per group) as depicted below, and the choice of dose

regimen for the extract is that the low dose is for establishing baseline response, the medium dose is assessing therapeutic or effective dose and the high dose is for determining toxicity and maximum tolerated dose (MTD)

- Group 1: Control (no treatment)  
 Group 2: MSU-induced arthritis + allopurinol tablet (5 mg/kg of body weight)  
 Group 3: MSU-induced arthritis + low-dose *C. afer* extract (100 mg/kg body weight)  
 Group 4: MSU-induced arthritis + medium-dose *C. afer* extract (200 mg/kg body weight)  
 Group 5: MSU-induced arthritis + high-dose *C. afer* extract (400 mg/kg body weight)

Treatment was administered orally once daily for 7 days, starting 24 hours after MSU injection.

## 2.7 Assessment of Serum Uric Acid, TNF- $\alpha$ , and IL-6 Levels

### 2.7.1 Collection of blood

At the end of the treatment period, blood samples were collected from the rats via cardiac puncture under anaesthesia into the sample blood bottles.

### 2.7.2 Determination of serum uric acid

*Serum uric acid* level was measured by using the enzymatic colourimetric method of Fossate et al. [27] with slight modification done by Hussain et al. [28].

### 2.7.3 Determination of serum TNF- $\alpha$ and IL-6 levels

The serum level of *TNF- $\alpha$*  was measured using ELISA kits as reported by Acar et al. [29]. In this method, blood samples collected in the plain tubes were centrifuged for around 10 min at  $\times 1500$  g which was followed by serum extraction. *TNF- $\alpha$*  levels were estimated by using a human "ELISA kit (Diacclone, Besancon Cedex, France) following the manufacturer's protocol. Similarly, the serum level of IL-6 was measured using ELISA kits as reported by Saxena et al. [30]. *IL-6* levels were determined from the serum samples extracted by centrifuging the blood collected in the plain vial at  $\times 1500$  g for 10 min by using a commercially available human *IL-16* ELISA kit from Krishgen Biosystems (United Kingdom) following the manufacturer's instructions.

## 2.8 Statistical Analysis

Data were analyzed using one-way ANOVA followed by Tukey's post hoc test to compare the means between groups, a *p*-value of  $<0.05$  was considered statistically significant. Descriptive statistics (mean  $\pm$  standard deviation) were used to summarize the data.

## 3. RESULTS

### 3.1 Phytochemical Analysis

The methanolic stem extracts of *C. afer* were analyzed for their phytochemical content. Qualitative analysis revealed the presence of flavonoids, phenols, tannins, saponins, and alkaloids, while anthraquinones, cardiac glycosides, terpenoids, phlobatannins, and cardenolides were not detected (Table 1). Quantitative analysis showed concentrations of flavonoids ( $0.774 \pm 0.036$  mg GAE/g), alkaloids ( $0.368 \pm 0.02$  mg GAE/g), and phenols ( $0.29 \pm 0.01$  mg GAE/g), along with lower levels of anthraquinones, cardiac glycosides, and terpenoids (Table 2).

**Table 1. Qualitative phytochemical screening results of methanolic stem extracts of *C. afer***

Phytochemical	Result
Flavonoids	+++
Tannins	-
Saponins	-
Alkaloids	+
Phenols	+++
Anthraquinones	++
Cardiac Glycosides	+
Terpenoids	++
Cardenolides	-
Terpenoids	-
Phlobatannins	-

**Key:** +: Present in trace amounts  
 ++: Present in moderate amounts  
 +++: Present in high amounts

**Table 2. Quantitative determination of the phytochemical content of methanolic stem extracts of *C. afer***

Phytochemical	Concentration (mg GAE/g)
Flavonoids	$0.774 \pm 0.036$
Alkaloids	$0.368 \pm 0.02$
Phenols	$0.29 \pm 0.01$
Anthraquinones	$0.012 \pm 0.003$
Cardiac glucosides	$0.011 \pm 0.002$
Terpenoids	$0.17 \pm 0.04$

### 3.2 Antioxidant Activity of Methanol Extract of *C. afer*

Table 3 presents the quantitative determination of antioxidant activity and the IC<sub>50</sub> values for the methanol extract of *C. afer* and standard antioxidants (gallic acid and ascorbic acid). The antioxidant activity of the methanol extract of *C. afer* was assessed across different concentrations, demonstrating a dose-dependent increase in antioxidant activity. At 100 µg/ml, the extract exhibited 67.14% antioxidant activity, which increased to 74.85% at 400 µg/mL. The IC<sub>50</sub> value of 92.33 µg/mL indicates the concentration required to inhibit 50% of free radical activity, suggesting a moderate antioxidant potential compared to the standards used.

Gallic Acid, a known potent antioxidant, showed higher activity across all concentrations with an IC<sub>50</sub> value of 37.92 µg/ml, reflecting its superior antioxidant capacity relative to the methanol extract of *C. afer*. Ascorbic Acid also displayed significant antioxidant activity with an IC<sub>50</sub> value of 51.40 µg/ml. Overall, while the methanol extract of *C. afer* demonstrates promising antioxidant activity, its effectiveness is lower compared to the standard antioxidants used in this study.

### 3.3 Serum Uric Acid, TNF-α Levels, and IL-6 Levels

The results of the effects of methanolic stem extracts of *C. afer* on serum uric acid, TNF-α, and IL-6 Levels are presented in Table 4.

**Table 3. Quantitative determination of antioxidant activity and IC<sub>50</sub> of methanol extract of *Costus afer* and standards**

Sample Extract	Concentration (µg/ml)	Percentage Antioxidant Activity (%)	IC <sub>50</sub> (µg/ml)
Methanol Extract of <i>C. afer</i>			<b>92.33</b>
	100	67.14 ± 1.58	
	200	68.67 ± 0.26	
	300	72.29 ± 0.26	
	400	74.85 ± 0.25	
Gallic Acid			<b>37.92</b>
	100	78.36 ± 1.05	
	200	82.24 ± 0.66	
	300	84.73 ± 0.66	
	400	86.67 ± 1.18	
Ascorbic Acid			<b>51.40</b>
	100	71.87 ± 2.21	
	200	76.43 ± 3.03	
	300	78.77 ± 2.91	
	400	88.97 ± 1.11	

**Table 4. Effects of methanolic stem extracts of *Costus afer* on serum uric acid, TNF-α, and IL-6 levels**

Parameter	Control Group	Untreated MSU Group	Low Dose (100 mg/kg)	Medium Dose (200 mg/kg)	High Dose (400 mg/kg)
Serum Uric Acid (mg/dL)	5.26 ± 0.45	9.32 ± 0.50**	7.58 ± 0.40*	6.32 ± 0.30**	4.78 ± 0.35***
TNF-α (pg/mL)	15.7 ± 1.2	28.4 ± 1.6**	22.5 ± 1.3*	17.8 ± 1.2**	12.4 ± 1.1***
IL-6 (pg/mL)	32.1 ± 2.0	58.9 ± 3.1**	48.2 ± 2.8*	38.6 ± 2.4**	28.1 ± 2.2***

\*Significant difference from untreated MSU group ( $p < 0.05$ )

\*\*Significant difference from the control group ( $p < 0.05$ )

\*\*\*Highly significant difference from the control group ( $p < 0.01$ )

### 3.4 Serum Uric Acid

The results of the effect of methanolic stem extracts of *Costus afer* on serum uric acid levels are presented in Table 4. The untreated MSU group exhibited significantly elevated *serum uric acid* levels ( $9.32 \pm 0.50$  mg/dL) compared to the control group ( $5.26 \pm 0.45$  mg/dL), indicating increased uric acid levels due to the condition being studied. Treatment with the low dose (100 mg/kg) of the extracts reduced serum uric acid to  $7.58 \pm 0.40$  mg/dL. The medium dose (200 mg/kg) resulted in a further decrease to  $6.32 \pm 0.30$  mg/dL, while the high dose (400 mg/kg) achieved the most substantial reduction, lowering serum uric acid to  $4.78 \pm 0.35$  mg/dL. This high dose resulted in levels significantly ( $p < 0.05$ ) below those of the control group, demonstrating a potent effect of the extracts in reducing *serum uric acid*.

### 3.5 Serum *TNF- $\alpha$* Levels

The results of the effect of methanolic stem extracts of *C. afer* on *TNF- $\alpha$*  levels are also detailed in Table 4. The untreated MSU group showed significantly higher *TNF- $\alpha$*  levels ( $28.4 \pm 1.6$  pg/mL) compared to the control group ( $15.7 \pm 1.2$  pg/mL), indicating increased inflammation. Treatment with the low dose (100 mg/kg) of the extracts reduced *TNF- $\alpha$*  to  $22.5 \pm 1.3$  pg/mL. The medium dose (200 mg/kg) further decreased *TNF- $\alpha$*  to  $17.8 \pm 1.2$  pg/mL, while the high dose (400 mg/kg) led to the most significant reduction, with *TNF- $\alpha$*  levels dropping to  $12.4 \pm 1.1$  pg/mL. This high dose resulted in *TNF- $\alpha$*  levels significantly ( $p < 0.05$ ) lower than those of the control group, suggesting a strong anti-inflammatory effect of the extracts.

### 3.6 Serum *IL-6* Levels

The effect of methanolic stem extracts of *C. afer* on *IL-6* levels, as presented in Table 4, showed that the untreated MSU group had elevated *IL-6* levels ( $58.9 \pm 3.1$  pg/mL) compared to the control group ( $32.1 \pm 2.0$  pg/mL). Treatment with the low dose (100 mg/kg) reduced *IL-6* to  $48.2 \pm 2.8$  pg/mL. The medium dose (200 mg/kg) further decreased *IL-6* to  $38.6 \pm 2.4$  pg/mL, and the high dose (400 mg/kg) achieved the most significant reduction, lowering *IL-6* to  $28.1 \pm 2.2$  pg/mL. This high dose resulted in *IL-6* levels significantly ( $p < 0.05$ ) lower than those of the control group, highlighting the effectiveness of the extracts in reducing inflammatory cytokine levels.

## 4. DISCUSSION

The study aimed to evaluate the therapeutic potential of methanolic stem extracts of *C. afer* in modulating serum uric acid, *TNF- $\alpha$* , and *IL-6* levels, which are key biomarkers in gouty arthritis and other inflammatory and metabolic disorders. The results demonstrate a notable dose-dependent effect of the extract on these parameters, highlighting its potential as a therapeutic agent.

The antioxidant activity of the methanol extract of *C. afer* was assessed, showing a dose-dependent increase with percentages ranging from 67.14% at 100  $\mu$ g/ml to 74.85% at 400  $\mu$ g/ml. The  $IC_{50}$  value (92.33  $\mu$ g/ml) of the extract indicates its moderate antioxidant potential compared to standards like Gallic Acid and Ascorbic Acid, which exhibited  $IC_{50}$  values of 37.92  $\mu$ g/ml and 51.40  $\mu$ g/ml, respectively. These findings suggest that while *C. afer* has significant antioxidant activity, it is less potent than the standard antioxidants. This result is consistent with other studies on *C. afer*, which have reported moderate antioxidant properties due to its phytochemical content, including flavonoids and phenolic compounds [22]. Antioxidants are implicated in decreasing morbidity and mortality arising from oxidative stress [31].

The significant reduction in serum uric acid levels observed with *C. afer* extracts supports its potential in managing hyperuricemia [32]. This finding aligns with research on other plant-based interventions such as *Hibiscus sabdariffa*, and *Aster glehni* which have demonstrated similar uric acid-lowering effects attributed to their ability to inhibit xanthine oxidase, an enzyme involved in uric acid production [33,34]. The high dose of *C. afer* extracts achieved a reduction in uric acid levels below the control, suggesting a potent effect comparable to or exceeding that observed with other plant extracts known for their uric acid-lowering properties.

The observed reduction in *TNF- $\alpha$*  levels with *C. afer* extracts highlights its potential anti-inflammatory properties. Previous research supports this finding, with various plant extracts such as *Curcuma longa* (turmeric) and *Zingiber officinale* (ginger) showing reductions in *TNF- $\alpha$*  levels in inflammatory models [35,36]. The high dose of *C. afer*

extracts significantly reduced *TNF- $\alpha$*  levels, indicating its effectiveness in modulating inflammation, consistent with findings from other studies on anti-inflammatory plant extracts.

Similarly, the reduction in *IL-6* levels observed with *C. afer* extracts is consistent with research on other anti-inflammatory agents. For example, *Panax ginseng* and *Andrographis paniculata* have been reported to reduce *IL-6* levels in animal models of inflammation [37]. The high dose of *C. afer* extracts achieved a significant reduction in *IL-6* levels, suggesting a strong anti-inflammatory action comparable to other plant-based therapies.

The anti-inflammatory property of *C. afer* may be attributed to its high content of flavonoids. Many studies have demonstrated that flavonoids can reduce cytokine expression and secretion, providing therapeutic benefits in inflammation-associated diseases as cytokine modulators [38,39]. Studies by Panche et al. [40] and Farraz et al. [41] opined that the anti-inflammatory effects of flavonoids are fundamentally due to their adeptness in modulating numerous biochemical pathways related to inflammation. They can inhibit the formation of pro-inflammatory cytokines for example *IL-1 $\beta$*  and *TNF- $\alpha$* , and enzymes like *COX-2*. Furthermore, flavonoids can subdue the stimulation of transcription factors like *NF- $\kappa$ B* and *STAT3*, which play influential roles in the inflammatory response. Certain flavonoids similarly interfere with the *NLRP3* inflammasome, a key protein complex involved in the inflammatory process. The phytochemical screening and quantitative analysis of *C. afer* support this hypothesis, suggesting that the observed anti-inflammatory effects may be mediated through these flavonoids and other active compounds present in the extract. This implies that the methanol extract of *C. afer* demonstrates notable antioxidant and anti-inflammatory properties, with significant effects on serum uric acid, *TNF- $\alpha$* , and *IL-6* levels. These findings highlight the potential of *C. afer* as a therapeutic agent for managing conditions such as gouty arthritis, though further research is needed to fully elucidate its mechanisms and optimize its therapeutic applications

## 5. CONCLUSION

The study successfully demonstrated that methanolic stem extracts of *C.s afer* exert

significant effects on serum uric acid, *TNF- $\alpha$* , and *IL-6* levels in a dose-dependent manner. The high dose of the extract achieved the most pronounced reductions in these biomarkers, suggesting a potent therapeutic potential. The reduction in serum uric acid levels indicates that *C. afer* may be effective in managing conditions associated with hyperuricemia, such as gout. The significant decrease in *TNF- $\alpha$*  and *IL-6* levels highlights the] potential anti-inflammatory properties of the extract, which could be beneficial in treating inflammatory disorders like gouty arthritis. Overall, the findings support the traditional use of *C. afer* and provide scientific validation for its therapeutic applications in managing inflammatory and metabolic conditions.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## CONSENT AND ETHICAL APPROVAL

It is not applicable.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Wu M, Tian Y, Wang Q, Guo C. Gout: A disease involved with complicated immunoinflammatory responses: A narrative review. *Clinical Rheumatology*. 2020;39(10):2849-2859.
2. Cha Y, Lee J, Choy W, Lee JS, Lee HH, Chae DS. Pathophysiology and treatment of gout arthritis; including gout arthritis of hip joint: a literature review. *Hip & Pelvis*. 2024;36(1):1.
3. Yokose C, Chen M, Berhanu A, Pillinger MH, Krasnokutsky S. Gout and osteoarthritis: associations, pathophysiology, and therapeutic implications. *Current rheumatology reports*. 2016;18:1-8.
4. Galozzi P, Bindoli S, Luisetto R, Sfriso P, Ramonda R, Scanu A, Oliviero F.

- Regulation of crystal induced inflammation: current understandings and clinical implications. *Expert Review of Clinical Immunology*. 2021;17(7):773-787.
5. Saribal D, Hocaoglu-Emre FS, Erdogan S, Bahtiyar N, Caglar Okur S, Mert M. Inflammatory cytokines IL-6 and TNF- $\alpha$  in patients with hip fracture. *Osteoporosis International*. 2019;30:1025-1031.
  6. Silva LB, dos Santos Neto AP, Maia SM, dos Santos Guimarães C, Quidute IL, Carvalho ADA, Leão JC. The role of TNF- $\alpha$  as a proinflammatory cytokine in pathological processes. *The Open Dentistry Journal*. 2019;13(1).
  7. Kany S, Vollrath JT, Relja B. Cytokines in inflammatory disease. *International journal of molecular sciences*. 2019;20(23):6008.
  8. Idrus HH, Budu HM. Biological effects of tumor necrosis factor alpha (TNF- $\alpha$ ) in systemic inflammation. *International journal dental and medical sciences research*. 2020;3(3):7-15.
  9. Roe K. An inflammation classification system using cytokine parameters. *Scandinavian journal of immunology*. 2021;93(2):e12970.
  10. Hirano T. IL-6 in inflammation, autoimmunity and cancer. *International immunology*. 2021;33(3):127-148.
  11. Narang RK, Dalbeth N. Pathophysiology of gout. In *Seminars in nephrology*. WB Saunders. 2020;40(6):550-563.
  12. Zhang T, Wang G, Zheng J, Li S, Xu, J. Profile of serum cytokine concentrations in patients with gouty arthritis. *Journal of International Medical Research*. 2021;49(11):03000605211055618.
  13. Shukla R, Pandey V, Vadnere GP, Lodhi S. Role of flavonoids in management of inflammatory disorders. In *Bioactive food as dietary interventions for arthritis and related inflammatory diseases*. Academic Press. 2019;293-322.
  14. Kpomah ED, Monday DA, Kpomah B. GCMS analysis of leaves and seeds of *Piper guineense* Schumach & Thoon. *African Scientist*. 2019;20(3):127-138
  15. Kpomah ED, Ogbogbo J, Kpomah B. Sub-acute toxicity studies of *Phyllanthus amarus* on haematological parameters and some plasma enzyme activities in mice. *International Journal of Basic Science and Technology*. 2017;3(1):53-58.
  16. Kpomah B, Egboh SHO, Agbaire PO, Kpomah ED. Spectroscopic characterization, antimicrobial and toxicological properties of derivatised thiosemicarbazone transition metal complexes” *Saudi. Journal of Medical and Pharmaceutical Sciences*. 2016;2(12):318-325
  17. Kpomah B, Kpomah ED. Toxicological assessment of acetone thiosemicarbazone metal complexes on body weight, biochemical parameters and liver histology of wistar rats. *Journal of Pharmaceutical and Applied Chemistry*. 2017;3(3):215-224.
  18. Kpomah ED, Kpomah B, Arhoghro EM. Histomorphological and biochemical changes induced in male wistar rats by chronic oral doses of *Piper guineense* Schumach. & Thonn. *Nigerian Journal of Pharmaceutical and Applied Science Research*. 2018;7(1):44-51.
  19. Kpomah ED, Arhoghro EM. Effects of doses of *Bryophyllum pinnatum* and glibenclamide on serum glucose and lipid profile in alloxan-diabetic rats. *Indian Journal of Drugs and Disease*. 2012;1(5):124-128
  20. Uchegbu R, Akalazu J, Ibe C, Ahuchaogu A, Amadikwa C. Chemical composition of the stem extract of *Costus afer* (Bush Cane) and its antimicrobial activity. *British Journal of Pharmaceutical Research*. 2016;10(5), 1-9.
  21. Anyasor GN, Onajobi F, Osilesi O, Adebawo O, Oboutor EM. Anti-inflammatory and antioxidant activities of *Costus afer* Ker Gawl. hexane leaf fraction in arthritic rat models. *Journal of ethnopharmacology*. 2014;155(1):543-551
  22. Anyasor GN, Onajobi F, Osilesi O, Adebawo O, Oboutor EM. Anti-inflammatory and antioxidant activities of *Costus afer* Ker Gawl. hexane leaf fraction in arthritic rat models, *Journal of Ethnopharmacology*. 2014;155(1):543-551.  
Available:<https://doi.org/10.1016/j.jep.2014.05.057>
  23. Kpomah ED, Onyeike EN, Kpomah B. Evaluation of some elemental, bioactive



- compounds and proximate composition of three commonly used herbal plants in the Niger Delta Region of Nigeria. *Chemistry Research Journal*. 2018; 3(2):12-21.
24. Wu Y, Wang X, Liu P, Niu Q, Wu Q. Quantitative determination of anthraquinones and resveratrol in *Polygonum Cillinerve* (Nakai) Ohwi by HPLC-PAD. *Journal of AOAC International*. 2017;100(1):25-29.
25. Ikeda Y, Fujii Y, Umemura M, Hatakeyama T, Morita M, Yamazaki, M. Quantitative determination of cardiac glycosides in *Digitalis lanata* leaves by reversed-phase thin-layer chromatography. *Journal of Chromatography A*. 1996;746(2): 255-260.
26. Malik SK. Qualitative and quantitative estimation of terpenoid contents in some important plants of Punjab, Pakistan. *Pakistan Journal of Science*. 2017;69(2): 150-154
27. Fossati P, Prencipe L, Berti G. Use of 3, 5-dichloro-2-hydroxybenzenesulfonic acid/4-aminophenazone chromogenic system in direct enzymic assay of uric acid in serum and urine. *Clinical chemistry*. 1980;26(2):227-231.
28. Hussain MS, Jahan N, Rashid MMO, Hossain MS, Chen U, Rahman N. Antihyperlipidemic screening and plasma uric acid reducing potential of *Momordica charantia* seeds on Swiss albino mice model. *Heliyon*. 2019;17:5(5):e01739. DOI: 10.1016/j.heliyon.2019.e01739
29. Acar L, Atalan N, Karagedik EH, Ergen A. Tumour necrosis factor-alpha and nuclear factor-kappa B gene variants in sepsis. *Balkan Medical Journal*. 2018;35: 30-35.
30. Saxena D, Dawson L, Gera R. Study of serum cytokines (interleukin-6 and tumor necrosis factor-alpha) in acute leukemias. *Journal of Radiation and Cancer Research*. 2022;13(2):74-80.
31. Arhoghro EM, Kpomah ED. Alanine Aminotransferase and aspartate aminotransaminase activities in wistar rats fed with *Musa paradisiaca* (Plantain) stem pulp in aluminium chloride induced hepatic oxidative stress. *Journal of Applied Science and Environmental Management*. 2022;26(6): 1057-1062
32. Kpomah ED Arhoghro EM. Positive stimulatory potentials of Coconut (*Cocos nucifera* L.) juice extract on *in-vivo* antioxidants, renal function and lipid profile of male wistar rats. *European Journal of Medicinal plants*. 2023;34(3): 45-54
33. Kuo CY, Kao ES, Chan KC, Lee HJ, Huang TF, Wang CJ. *Hibiscus sabdariffa* L. extracts reduce serum uric acid levels in oxonate-induced rats. *Journal of Functional Foods*. 2012;4(1):375-381.
34. Park JE, Yeom Z, Park KT, Han EH, Yu HJ, Kang HS, Lim YH. Hypouricemic effect of ethanol extract of *Aster glehni* leaves in potassium oxonate-induced hyperuricemic rats. *Clinical nutrition research*. 2018; 7(2):126.
35. Azeez TB, Lunghar J. Antiinflammatory effects of turmeric (*Curcuma longa*) and ginger (*Zingiber officinale*). *Inflammation and Natural Products*. 2021; 83-102.
36. Akinyemi AJ, Thomé GR, Morsch VM, Bottari NB, Baldissarelli J, de Oliveira LS, Schetinger MRC. Effect of ginger and turmeric rhizomes on inflammatory cytokines levels and enzyme activities of cholinergic and purinergic systems in hypertensive rats. *Planta medica*. 2016;82(07):612-620.
37. Widjowati R, Pratama RR, Sholikhah I, Jain SK. Herbal medicine and rheumatic disorders management and prevention. In *Herbal Medicine Phytochemistry: Applications and Trends* (pp. 723-762). Cham: Springer International Publishing; 2024.
38. Al-Khayri JM, Sahana GR, Nagella P, Joseph BV, Alessa FM, Al-Mssallem MQ. Flavonoids as potential anti-inflammatory molecules: A review. *Molecules*. 2022;27(9):2901.
39. Maleki SJ, Crespo JF, Cabanillas B. Anti-inflammatory effects of flavonoids. *Food chemistry*. 2019;299:125124.
40. Panche AN, Diwan AD, Chandra SR. Flavonoids: An overview. *Journal of Nutritional Science*. 2016;5:e47. DOI:10.1017/jns.2016.4
41. Ferraz CR, Carvalho TT, Manchope MF, Artero NA, Rasquel-Oliveira FS, Fattori V, Casagrande R, Verri WA Jr. Therapeutic potential of flavonoids in pain and inflammation: Mechanisms of action, pre-

clinical and clinical data, and  
pharmaceutical development. *Molecules*.  
2020;25(3):762.

Available:[https://doi.org/10.3390/molecules  
25030762](https://doi.org/10.3390/molecules25030762)

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/124422>