



Comparative Intraocular Pressure Study among Diabetic and Non-Diabetic Patients

Asadullah Jatoi ^{a*}, Noman Ahmed Shaikh ^a and Mona Liza Mahesar ^a

^a *Institute of Ophthalmology, Liaquat University of Medical & Health Sciences, Jamshoro, Sindh, Pakistan.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i53B33699

Editor(s):

(1) Dr. Debarshi Kar Mahapatra, Rashtrasant Tukadoji Maharaj Nagpur University, India.

Reviewers:

(1) Rajesh Jain, India.

(2) Rateesh Sareen, Santokba Durlabhji Memorial Hospital & Research Centre, India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/78002>

Original Research Article

Received 28 September 2021

Accepted 01 December 2021

Published 08 December 2021

ABSTRACT

Background: Diabetes mellitus is one of the major health issues in the world. Its microvascular complications contribute to ocular complications including increased intraocular pressure (IOP) which is a risk factor of glaucoma. Identification of factors responsible for glaucoma is a mainstay in the early detection and prevention of blindness.

Aim and Objectives: The objective of the study was to compare IOP among diabetic and non-diabetic patients and to assess the correlation between age and IOP.

Materials and Methods: The cross-sectional study was performed on 104 participants. Patients were divided into two groups based on diabetes (case, n = 52) and non-diabetes (control, n = 52). The detailed history and routine clinical investigations were performed. Three consecutive readings of IOP of the left and right eye were recorded separately using Goldmann applanation tonometer. Wilcoxon sign-rank test and Spearman-correlation test was used to find the difference between the IOP and correlation between age and IOP, respectively.

Results: A significantly higher IOP was observed in diabetic patients than non-diabetic patients. (0.0001). There was no sex difference in IOP was observed in diabetic and non-diabetic patients. Age of the patient was not correlated with IOP ($r = -0.02197219$, $P = 0.824$).

Conclusion: Diabetic patients are prone to higher IOP; therefore, diabetic patients should be regularly assessed for IOP for diagnosis glaucoma.

Keywords: Blindness; diabetes mellitus; glaucoma; intraocular pressure; open-angle glaucoma.

1. INTRODUCTION

Globally diabetes became a pandemic. It was estimated that in the year 2017, there were 451 million people with diabetes mellitus (DM) and most of the people were 40–60-year age. These numbers are expected to increase to 693 million by 2045 [1]. The prevalence of diabetes in Asian countries is high and it accounts for >60% of the global diabetic population [2]. In Pakistan, currently, 62 million people are affected with DM and these numbers are predicted to be increased to 79.4 million by the year 2030 [3]. It causes major health burden causing substantial financial loss because of higher rates of morbidity and mortality and health care expenditures [4].

DM is commonly associated with microvascular complications contributing to various ocular complications such as increased intraocular pressure (IOP) and subsequent glaucoma which is a common cause irreversible blindness [5-10]. Diabetics are more prone to have primary open-angle glaucoma than non-diabetics [11,12]. To be noted, glaucoma is estimated to affect 12 million Pakistanis: Accounting for 12.8% of total blindness and is considered to be the 3rd most common reason for blindness in the country [13]. IOP is the pressure exerted by the fluid inside the eye. It is an important ophthalmic physiological parameter important to understand the distribution and risk factors of IOP for the prevention and prognosis of glaucoma [14]. Various factors such as age, body mass index, blood pressure, and central corneal thickness are associated with IOP [15-17]. However, the results of the various studies were not entirely consistent, and the potential risk factors in their analysis were failed to account due to a lack of data [14]. Moreover, it remains ambiguous whether DM population has distinct distribution or risk factors for IOP, and the association of DM with glaucoma has still been controversial, despite this DM individuals are twice likely to develop glaucoma compared to non-diabetic individuals [18]. Therefore, data regarding IOP distribution and risk factors in DM individuals are required to produce the relationship between glaucoma and DM and plan effective strategies. The present study aimed to compare IOP among diabetic and non-diabetic patients and to assess the correlation between age and IOP.

2. MATERIALS AND METHODS

The cross-sectional study was conducted at the tertiary care center at Jamshoro and Hyderabad.

A total of 104 Types 2 diabetic and non-diabetic subjects of 40–60 years of age were included in the study and informed consent was obtained. Patients with Type 1 DM, gestational diabetes, intraocular tumors, and glaucoma were excluded from the study. Patients were divided into two groups based on diabetes (cases), and non-diabetes (control). Each group consisted of $n = 52$ patients. The detailed history of patients and routine clinical investigations were performed. Three consecutive readings of IOP were recorded by Goldmann applanation tonometer. The IOP of the left and the right eye was recorded separately after anesthetizing eye with xylocaine during outpatient department hours.

2.1 Statistical Analysis

Data were analyzed using R Studio V 1.2.5001 software. Continuous variables were expressed in mean \pm SD whereas, categorical variables were expressed as percentage and frequency. Wilcoxon sign-rank test was used to find the difference between the IOP and Spearman-correlation test was used to find the correlation between age and IOP. $P < 0.05$ was considered statistically significant.

3. RESULTS

The average age of the participants was 55.83 ± 5.87 years and the majority of the participants were female (58%). Both groups were similar in terms of age and sex. A significantly higher IOP was observed in diabetic patients than non-diabetic patients (0.0001) [Table 1]. A significantly higher IOP was observed in diabetic male and female patients than non-diabetic male and female patients. No significant difference was observed between the IOP of males and females of both groups [Table 2]. No correlation was observed between age and IOP ($r = -0.02197219$, $P = 0.824$) [Fig. 1].

4. DISCUSSION

In diabetic patients, the IOP was higher than non-diabetic patients ($P = 0.0001$). Among diabetic male and female patients, no sex difference was observed. Similarly, there was no correlation between age and IOP. The study showed significantly higher IOP in diabetic patients (15.96 ± 2.27 mmHg) than non-diabetic patients (13.84 ± 2.94 mmHg) ($P = 0.0001$). These findings are following the previous reports [3,7,19]. The exact mechanism of increased IOP

Table 1. Distribution of demographical variables and IOP according to groups

Variable	Control	Cases	P-value
Age (years)	56.96±4.89	54.71±6.58	0.1694
Sex			
Female	29	29	1.000
Male	23	23	
IOP (mmHg)	13.84±2.94	15.96±2.27	0.0001***

***P<0.001, IOP: Intraocular pressure

Table 2. Distribution of IOP according to sex

Sex	IOP		P-value
	Control	Cases	
Male	14.19±2.57	15.96±2.47	0.024*
Female	13.53±2.68	15.41±1.94	0.007**
P-value	0.4498	0.3368	

*P<0.05, **P<0.01, IOP: Intraocular pressure




Fig. 1. Scatter plot between age and intraocular pressure

in DM patients is not known. However, in vitro study suggested that increased deposition of fibronectin in the extracellular matrix of tubercular meshwork blocks the aqueous outflow which leads to decreased aqueous drainage hence, a rise in IOP [20]. In this study, no sex difference was observed. However, studies have reported that hormonal differences may play a role in the sex difference of IOP, resulting in higher IOP in female than male participants [14,21,22]. The difference in the reports may be due to the ethnic/racial difference, method of IOP assessment, or lifestyle of the population. In this series, no correlation was observed between age and IOP ($r = -0.02197219$, $P = 0.824$). However, Vidhya et al. in their study suggested that increase in mean IOP with each decade of life.[19] Despite this, there is no general agreement on the association between IOP and age in the literature. Studies reporting increased, decreased, and no association of IOP with age

[23-25]. The difference in the results may be due to different populations, or a nonmonotonic relationship between age and IOP such as different studies with different aged participants [26]. The present study showed higher IOP in diabetic compared to non-diabetic patients. These findings suggest that diabetic patients should undergo routine IOP assessment to avoid chances of glaucoma. The limitation of the study was the small sample size, the parameters such as blood pressure, body mass index, and obesity were not assessed. A study with a large sample size including all variables is the further recommendation for the confirmation of the present study findings.

5. CONCLUSION

The conclusion of the study suggested an increased IOP in diabetes patients and no sex difference in IOP of diabetic and non-diabetic

patients. Furthermore, no correlation was found between age and IOP. Since IOP is a known risk factor for glaucoma in diabetic patients, this would suggest that diabetics be monitored regularly for IOP for early diagnosis of glaucoma in the susceptible patients.

CONSENT

As per international standard or university standard, patient's written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Cho NH, Shaw JE, Karuranga S, Huang Y, da Rocha Fernandes JD, Ohlrogge AW, et al. IDF diabetes atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res Clin Pract.* 2018;138:271-81.
2. Malik VS, Willett WC, Hu FB. Global obesity: Trends, risk factors and policy implications. *Nat Rev Endocrinol.* 2013;9:13-27.
3. Baisakhiya S, Garg P, Singh S. Association between glycemic control and intraocular pressure in patients with Type II diabetes mellitus. *Natl J Physiol Pharm Pharmacol.* 2017;7:43-6.
4. Bommer C, Sagalova V, Heeseemann E, Manne-Goehler J, Atun R, Bärnighausen T, et al. Global economic burden of diabetes in adults: Projections from 2015 to 2030. *Diabetes Care.* 2018;41:963-70.
5. Dielemans I, de Jong PT, Stolk R, Vingerling JR, Grobbee DE, Hofman A. Primary open-angle glaucoma, intraocular pressure, and diabetes mellitus in the general elderly population: The Rotterdam study. *Ophthalmology.* 1996;103:1271-5.
6. Sahin A, Bayer A, Özge G, Mumcuoglu T. Corneal biomechanical changes in diabetes mellitus and their influence on intraocular pressure measurements. *Invest Ophthalmol Vis Sci.* 2009;50:4597-604.
7. Pimentel LG, Gracitelli CP, da Silva LS, Souza AK, Prata TS. Association between glucose levels and intraocular pressure: Pre-and postprandial analysis in diabetic and nondiabetic patients. *J Ophthalmol.* 2015;2015:832058.
8. Tan GS, Wong TY, Fong CW, Aung T. Diabetes, metabolic abnormalities, and glaucoma: The Singapore Malay eye study. *Arch Ophthalmol.* 2009;127:1354-61.
9. Zhao D, Cho J, Kim MH, Friedman DS, Guallar E. Diabetes, fasting glucose, and the risk of glaucoma: A meta-analysis. *Ophthalmology.* 2015;122:72-8.
10. Luo XY, Tan NY, Chee ML, Shi Y, Tham YC, Wong TY, et al. Direct and indirect associations between diabetes and intraocular pressure: The Singapore epidemiology of eye diseases study. *Invest Ophthalmol Vis Sci.* 2018;59:2205-11.
11. Arora VK, Prasad VN. Intraocular pressure and diabetes-a correlative study. *Indian J Ophthalmol.* 1989;37:10-2.
12. Singh M, Heong SC. Postural behaviour of intraocular pressure in diabetics. *Br J Ophthalmol.* 1986;70:456-9.
13. Krishnaiah S, Kovai V, Srinivas M, Bindiganavale RS, Rao GN, Thomas R. Awareness of glaucoma in the rural population of Southern India. *Indian J Ophthalmol.* 2005;53:205-8.
14. Cui Y, Yang X, Zhang G, Guo H, Zhang M, Zhang L, et al. Intraocular pressure in general and diabetic populations from Southern China: The Dongguan eye study. *Invest Ophthalmol Vis Sci.* 2019;60:761-9.
15. Hennis A, Wu SY, Nemesure B, Leske MC, Barbados Eye Studies Group. Hypertension, diabetes, and longitudinal changes in intraocular pressure. *Ophthalmology.* 2003;110:908.
16. Nomura H, Shimokata H, Ando F, Miyake Y, Kuzuya F. Age-related changes in intraocular pressure in a large Japanese population: A cross-sectional and longitudinal study. *Ophthalmology.* 1999;106:2016-22.
17. Wang D, Huang W, Li Y, Zheng Y, Foster PJ, Congdon N, et al. Intraocular pressure, central corneal thickness, and glaucoma in Chinese adults: The Liwan eye study. *Am J Ophthalmol.* 2011;152:454-62.
18. Zhao YX, Chen XW. Diabetes and risk of glaucoma: A systematic review and a meta-analysis of prospective cohort studies. *Int J Ophthalmol.* 2017;10:1430-5.

19. Vidhya NP, Das S, Priyadarshini R, Subashini M, Mahadevan K. A comparative study on the intraocular pressure among diabetic and non-diabetic patients. *Indian J Clin Exp Ophthalmol.* 2016;2:378-80.
20. Sato T, Roy S. Effect of high glucose on fibronectin expression and cell proliferation in trabecular meshwork cells. *Invest Ophthalmol Vis Sci.* 2002;43:170-5.
21. Astrom S, Stenlund H, Linden C. Intraocular pressure changes over 21 years-a longitudinal age-cohort study in Northern Sweden. *Acta Ophthalmol.* 2014;92:417-20.
22. Han X, Niu Y, Guo X, Hu Y, Yan W, He M, et al. Age- related changes of intraocular pressure in elderly people in Southern China: Lingtou eye cohort study. *PLoS One.* 2016;11:e0151766.
23. Lin HY, Hsu WM, Chou P, Liu CJ, Chou JC, Tsai SY, et al. Intraocular pressure measured with a noncontact tonometer in an elderly Chinese population: The Shihpai eye study. *Arch Ophthalmol.* 2005; 123:381-6.
24. Memarzadeh F, Ying-Lai M, Azen SP, Varma R, Los Angeles Latino Eye Study Group. Associations with intraocular pressure in Latinos: The Los Angeles Latino eye study. *Am J Ophthalmol.* 2008;146:69-76.
25. Klein BE, Klein R, Linton KL. Intraocular pressure in an American community. The Beaver Dam eye study. *Invest Ophthalmol Vis Sci.* 1992;33:2224-8.
26. Khawaja AP, Springelkamp H, Creuzot-Garcher C, Delcourt C, Hofman A, Höhn R, et al. Associations with intraocular pressure across Europe: The European eye epidemiology (E3) consortium. *Eur J Epidemiol.* 2016;31:1101-11.

© 2021 Jatoi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78002>*