



A Study on the Identification and Comparison of Functional Components in Different Species of Passion Fruit Vine Leaves

Poornima Jeyasekaran^{a*} and M. Deepa^{a#}

^a Department of Food Science and Nutrition, Periyar University, Salem, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i58B34230

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/79819>

Original Research Article

Received 10 October 2021

Accepted 14 December 2021

Published 16 December 2021

ABSTRACT

Functional components are found abundantly in plants which are non-nutritive and bio active compounds that prevents the onset of degenerative diseases and protects the body by maintaining the health. The objective of this study is to identify the chemical components of different varieties of passion fruit leaves (*Passiflora edulis f. flavicarpa*, *Passiflora edulis f. edulis* and *Passiflora quadrangularis*) available wildy in the hills of Tamil Nadu, India. Young tender passion fruit leaves can be used as a raw leafy green as they contain vitamin A and niacin. The dried leaves are used for calming teas and herbal remedies. The functional components were identified using Fourier Transform Infrared Spectroscopy FTIR(ATR) spectrophotometer. The leaves contain polyphenols, triterpenes, carotenoids, polysaccharides, amino acids and flavonoids which have a lot of beneficial effect on human health especially in preventing degenerative diseases. This paves way to use these amazing abundant greens in clinical studies as ayurvedic preparations and treat illness naturally.

Keywords: *Functional compounds; Passiflora edulis f. flavicarpa; Passiflora edulis f. edulis and Passiflora quadrangularis; FTIR(ATR).*

^{*}Research Scholar (Part-time);

[#]Assistant Professor;

^{*}Corresponding author: E-mail: poornimajeyasekaran@yahoo.co.in;

1. INTRODUCTION

Plants are an integral part of all the living organisms of earth because they provide food, clothing, shade and shelter. They also provide clean air and natural medicines to treat ailments. [1]. Many scriptures state that from the day life begins plants are used as medicines. Countries such as Egyptian, Assyrian, Chinese, and Indian, have long histories of using medicinal plants [2].

Plants are able to produce a large number of diverse bioactive compounds [3]. Plant kingdom mainly contains beneficial phytochemicals may complement the needs of the human body by acting as natural antioxidants [4]. These antioxidants compounds are secondary plant metabolites. Phenolic derivatives are the main phytochemicals present in the most of the species. Several clinical researches showed that antioxidant and their phytochemicals can prevent some chronic diseases [5,6].

Passion fruit leaves, grow on a fast-climbing vine that can spread 4-6 meters a year and are members of the Passifloraceae family. There are four hundred species within the Passifloraceae family. Nowadays the passion fruit plant which was mainly grown for its fruits is also growing in popularity as an additional crop. The leaves can be used as a culinary ingredient as well. Passion fruit vines are native from Brazil, but now it is cultivated in all parts of the world for the fruits and also as ornamental plants[7].

Traditionally *P. edulis* leaves are used in the treatment of insomnia and known to produce a restful sleep without any narcotic hangover. The leaves are reported to contain a bitter principle maracugine, resins, acids and tannin exceptionally rich in ascorbic acid. It is also used to treat epilepsy, ulcers and haemorrhoids [8]. The tea prepared by the infusion of these leaves has been recognized for its anti-inflammatory potential and also there was also an increase in colonic bacteria; therefore, it was associated with increase in SCFA production which promote different aspects of gut microbiome [9]. Studies proved that introduction of infusion of *P. edulis* leaves in tea could contribute to prevent damage by reactive species [10]. The presence of fructo-oligosaccharides in passion fruit by-products shows a comparable stimulation of folate production was found by the selected bacterial strains [11].

2. METHODOLOGY

The aim of the study is to identify the functional compounds in three different species of passion fruit leaves using FTIR (ATR) technique. The passion fruit vines of three different species *Passiflora edulis* f. *flavicarpa*, *Passiflora edulis* f. *edulis* and *Passiflora quadrangularis* were collected from Horticulture research Station in Thandiyan kudisai, Perumbarai, Dindigul district, Tamil Nadu, India in February 2020 at the evening time. The collection place was about 50 km away from the residence. Soon after collection, the samples were washed thoroughly and then dried at room temperature 30 degree Celsius and under shade for one week till completely dried as mentioned in Fig. 1. The examined moisture content was 10 percent. Then, the extract was prepared from the coarse powder with methanolic extraction method. The solid to solvent ratio is 1:10 and the extraction time was 24 hours. It was transferred to the separatory funnel and the extract was stored in sterile vials and taken to the central Instrumentation laboratory of the American college campus for FTIR (ATR) spectrophotometric analysis.




Fig. 1. Flowchart of FTIR Spectroscopy analysis process

2.1 Fourier Transformed Infrared Spectroscopy (FTIR)

The functional groups of *Passiflora* species were recorded using Fourier Transform Infrared Spectroscopy using JASCO spectrophotometer (FTIR 4600) ATR techniques. Infrared spectra were recorded between ranges 4000 and 450 cm^{-1} . Attenuated total reflection is developed in order to enhance the surface sensitivity since IR spectroscopy is a bulk method [12].

2.2 Fourier Transformed Infrared Spectroscopy (FTIR)

The functional groups of *Passiflora* species were recorded using Fourier Transform Infrared Spectroscopy using JASCO spectrophotometer (FTIR 4600) ATR techniques. Infrared spectra were recorded between ranges 4000 and 450 cm^{-1} . Attenuated total reflection is developed in order to enhance the surface sensitivity since IR spectroscopy is a bulk method [12].

3. RESULTS AND DISCUSSION

FTIR(ATR) analysis was done for the characterization of functional group present in the three different varieties of passion fruit vines namely *Passiflora edulis* f. *flavicarpa*, *Passiflora edulis* f. *edulis* and *Passiflora quadrangularis*. The interpretation of IR spectra helps to determine the functional groups present within organic

compounds. This can be determined by examining each spectrum for common IR stretches that are characteristics of organic functional groups. The current research reconfirms the presence of more than 110 phytochemical constituents identified from the different plant parts of *P. edulis* in which flavonoids and triterpenoids held the biggest share done by He et al., in 2020 [17].

Table 1 and Fig. 2 show the absorbance spectra recorded for *Passiflora edulis* f. *flavicarpa*.

“The strong peak frequency observed for *Passiflora edulis* f. *flavicarpa* 3346.85 cm^{-1} same to O-H extending and is between the standard frequencies (4000 – 3000 cm^{-1}). The peak frequency observed at 2926.45 cm^{-1} corresponds to the O-H stretching and the standard frequency is 3000-2500 cm^{-1} . The strong peak frequency observed at 1737.55 cm^{-1} corresponds to the C=O extending and the standard frequency is 2000-1650 cm^{-1} . The peak frequency observed at 1367.28 cm^{-1} corresponds to the O-H bending and the standard frequency is 1600-1300 cm^{-1} . The peak frequency observed at 1016.3 cm^{-1} corresponds to the C-F extending and the standard frequency is 1400-1000 cm^{-1} . Thus shows the possibility of flavonoids, phenols, alkaloids and amines as phytochemicals.

Table 2 and Fig. 3 show the absorbance spectra recorded for *Passiflora edulis* f. *edulis*.

Table 1. FTIR values of *Passiflora edulis* f. *flavicarpa* leaf

S.No	Frequency range (cm^{-1})	Absorption (cm^{-1})	Intensity	Group	Compound
1	4000 – 3000	3346.85	Strong broad	O-H Stretching	Alcohol
2	3000-2500	2926.45	Weak, broad	O-H Stretching	Alcohol
3	2000-1650	1737.55	Strong	C=O Stretching	Aldehyde
4	1600-1300	1367.28	Medium	O-H Bending	Alcohol
5	1250-1020	1215.9	Medium	C-N Stretching	Amine
6	1400-1000	1016.3	Strong	C-F Stretching	Fluoro compound

Table 2. FTIR values of *Passiflora edulis* f. *edulis* leaf

S. No	Frequency range (cm^{-1})	Absorption (cm^{-1})	Intensity	Group	Compound
1	4000-3000	3332.39	Strong broad	O-H Stretching	Alcohol
2	3000-2500	2945.73	Weak broad	O-H Stretching	Alcohol
3	3000-2500	2832.92	Weak broad	O-H Stretching	Alcohol
4	2000-1650	1737.55	Strong	C=O Stretching	Aldehyde
5	1400-1000	1368.25	Medium	C-H Bending	Alkane
7	1400-1000	1019.15	Strong sharp	C-F Stretching	Fluoro compound




Fig. 2. Functional components in *Passiflora edulis* f. flavicarpaleaf




Fig. 3. FTIR Spectra of *Passiflora edulisf. edulis* leaf

“The strong peak frequency observed for *Passiflora edulisf. edulis* 3332.39 cm⁻¹ same to O-H extending and is between the standard frequencies (4000 – 3000 cm⁻¹). The peak frequency observed at 2945.73 cm⁻¹ and 2832.92 cm⁻¹ corresponds to the O-H stretching and the standard frequency is 3000-2500 cm⁻¹. The strong peak frequency observed at 1737.55cm⁻¹ corresponds to the C=O extending and the standard frequency is 2000-1650 cm⁻¹. The peak frequency observed at 1368.25 cm⁻¹ corresponds to the C-H bending and the standard frequency is 1400-1000 cm⁻¹. The peak frequency observed at 1019.15 cm⁻¹ corresponds to the C-F extending and the standard frequency is 1400-1000 cm⁻¹. Thus shows the possibility of flavonoids, phenols, alkaloids and amines as phytochemicals.

Table 3 and Fig. 4 show the absorbance spectra recorded for *Passiflora quadrangularis*.

“The peak frequency observed for *Passiflora quadrangularis* 3302.5cm⁻¹ same to O-H extending and is between the standard frequencies (4000 – 3000 cm⁻¹). The strong and broad peak frequency observed at 2926.45 cm⁻¹ corresponds to the N-H stretching and the standard frequency is 4000-3000 cm⁻¹. The strong peak frequency observed at 1737.55cm⁻¹ corresponds to the C=O extending and the standard frequency is 2000-1650 cm⁻¹. The peak frequency observed at 1606.41cm⁻¹ corresponds to the C=Cstretching and the standard frequency is 1600-1300 cm⁻¹. The peak frequency observed at 1448.28 cm⁻¹ corresponds to the O-H bending and the standard frequency is 1400-1000 cm⁻¹. The peak frequency observed at 1369.21 cm⁻¹ corresponds to the C-F extending and the standard frequency is 1400-1000 cm⁻¹. The peak frequency observed at 1068.37cm⁻¹ corresponds to the S=O extending and the standard frequency is 1400-1000 cm⁻¹. The peak

frequency observed at 777.17 cm⁻¹ corresponds to the C-H bending and the standard frequency is 900-700 cm⁻¹. Thus shows the presence of the following probable phytochemicals: Alkaloids, carboxylic acids, flavonoids, ketones, tannins, poly phenols and sulfuraphanes.

The above results support the previous studies which states that apart from being a food item, a variety of pharmaceutical products based on ingredients have also been developed and used in folk medicine. The principal components of *P. edulis* include polyphenols, triterpenes, carotenoids, polysaccharides, amino acids, essential oils, and microelements [13,14].

Also, certain biogenic amines (catecholamines) which are produced by the decarboxylation of

amino acids or by the amination of aldehydes and ketones. Catecholamines include dopamine, serotonin, epinephrine, and norepinephrine and are reported to occur in many plants in considerable amounts [15].

The aqueous extract of *P. edulis* leaves showed a significant anti-inflammatory property on mice [16]. The general administration of *P. edulis* exhibited distinct anti-inflammatory actions which inhibits leukocyte influx to the pleural cavity and associated with marked blockade of myeloperoxidase, nitric oxide, TNF levels in the acute model of inflammation caused by intra pleural injection of mice. In one experiment, *P. edulis* was very effective in defeating the TNF levels than dexamethasone [10].

Table 3. FTIR values of *Passiflora quadrangularis* leaf

S.No	Frequency range (cm ⁻¹)	Absorption (cm ⁻¹)	Intensity	Group	Compound
1	4000-3000	3302.5	Strong broad	O-H Stretching	Alcohol
2	4000-3000	2926.45	Strong broad	N-H Stretching	Amine salt
3	2000-1650	1737.55	Strong	C=O Stretching	Esters
4	1600-1300	1606.41	Strong	C=C Stretching	Unsaturated ketone
5	1600-1300	1448.28	Medium	O-H Bending	Carboxylic acid
6	1400-1000	1369.21	Strong sharp	C-F Stretching	Fluoro compound
7	1400-1000	1068.37	Strong	S=O Stretching	Sulfoxide
8	900-700	777.17	Strong	C-H Bending	Disubstituted




Fig. 4. FTIR Spectra of *Passiflora quadrangularis* leaf

4. CONCLUSIONS

The three varieties of passion fruit leaves confirm the presence of different functional components which is a boon to human health maintenance and prevention of certain diseases. Though certain diseases were treated in the hospital other minor ailments such as headaches, stomach upset, colds and fevers are treated by medicinal plants as a home remedy. Passion fruit leaves can be used in both raw and cooked applications such as boiling, sautéing, and frying. Passion fruit leaves can also be cooked into soups, curries, sauté veggies, pasta, and quiches. They are similar in texture and flavour to spinach and can often be substituted in recipes for the green. Tea is the second most popular beverage in the world in addition to cooking, Passion fruit leaves can be boiled and made into a calming tea. The researcher recommend that passion fruit vines can be cultivated at home as grow your own greens and get a lot of health benefits over culinary uses.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENTS

The researcher acknowledges her heart felt gratitude to Dr. T. Thangaselvambai, Head of Horticulture Research Station, Thandiyankudisai, Perumbarai, Dindigul district, Tamil Nadu, India for contributing fresh Passion fruit vines for the study and Mr. N. Jeyasekaran and Mr. S. Jeyasudhan for transporting the fresh fruits to the laboratories. Sincere thanks to Dr. M. Davamani Christober, Principal & Secretary of the American college, Madurai for instrumentation facility. Thankful to Dr. Poongodi Vijayakumar and Dr. M. Deepa for their guidance.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ram JS. Genetic resources chromosome engineering & crop improvement, Taylor & Francis Group.2012;13.
2. Ramawat KG, Dass S, Meeta Mathur. Herbal drugs: Ethnomedicine to modern medicine chapter 2 the chemical diversity of bioactive molecules and therapeutic potential of medicinal plants. Springer, e-ISBN: 978-3-540-79116-4. 2008; 7-8.
3. Naczki M, Shahidi F. Phenolics in cereals, fruits and vegetables: Occurrence, extraction and analysis. J. Pharm. Biomed. Anal. 2006;41:1523–1542.
4. Boots AW, Haenen GR, Bast A. Health effects of quercetin: From antioxidant to nutraceutical. Eur. J. Pharmacol., 2008; 585:325–337.
5. Lebovka N, Vorobiev E, Chemat F. Enhancing extraction processes in the food industry. Taylor & Francis group, LLC.2012;268.
6. Zhimin XV, Howard LR. Analysis of antioxidant-rich phytochemical. John Wiley & Sons Ltd., 2012;1.
7. Sunitha M, Devaki K. Antioxidant activity of *Passiflora edulis* Sims leaves, Indian Journal of Pharmaceutical Sciences.2009; 71(3):310-311.
8. Relw S, Espig G. The cultivated plants of the tropic and subtropics. Rev ed. New Delhi: CBS Publishers and Distributors; 1991.
9. Juliana Kelly da Silva et al. Antioxidant activity of aqueous extract of passion fruit (*Passiflora edulis*) leaves: In vitro and in vivo study, Food Research International. 2013;53(2):882-890.
10. Montanher A, Zucolotto SM, Schenkel E, Frode T. Evidence of anti-inflammatory effects of *Passiflora edulis* in an inflammation model. J. Ethnopharmacol. 2007;109:281-288.
11. Albuquerque MAC, Bedani R, LeBlanc JG, Saad SMI. Passion fruit byproduct and fructooligosaccharides stimulate the growth and folate production by starter and probiotic cultures in fermented soymilk. International Journal of Food Microbiology.2017;261:35–41.

- Available:<https://doi.org/10.1016/j.ijfoodmicro.2017.09.001>.
12. AsepBayu Dani Nandiyanto, RosiOktiani, RistiRagadhita. How to read and interpret ftir spectroscope of organic material. Indonesian Journal of Science & Technology. 2019;4(1):97-118.
 13. He MJ, Zeng JY, Zhai L, Liu YG, Wu HC, Zhang RF, et al. Effect of in vitro simulated gastrointestinal digestion on polyphenol and polysaccharide content and their biological activities among 22fruit juices. Food Res. Int. 2017;102:156–162. DOI: 10.1016/j.foodres.2017.10.001.
 14. Yuan TZ, Kao CL, Li WJ, Li HT, ChenCY. Chemical constituents of leaves of *Passiflora edulis*. Chem. Nat. Compd. 2017;53:1165–1166. DOI: 10.1007/s10600-017-2227-5.
 15. Jiwan S. Sidhu, Tasleem A. Zafar. bioactive compounds in banana fruits and their health benefits. Food Quality and Safety, 2018;2(4):183–188. Available:<https://doi.org/10.1093/fqsafe/fyy019>.
 16. Vargas AJ, Geremias DS, Provensi G, Fornari PE, Reginatto FH, Gosmann G, Schenkel EP, Frode TS. *Passiflora alata* and *Passiflora edulis* spray dried aqueous extracts inhibit Inflammation in mouse model of pleurisy. Fitoterapia.2007;78: 112-119.
 17. Xirui He, Fei Luan, Yan Yang, Ze Wang, Zefeng Zhao, Jiacheng Fang, Min Wang, Manhua Zuo, Yongsheng Li. Phytochemistry and Pharmacology of *Passiflora edulis*. Front. Pharmacol. 2020;11:617. DOI: 10.3389/fphar.2020.00617

© 2021 Jeyasekaran and Deepa; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/79819>