



# Promising Combination Systemic Fungicides in Combating Basal Stem Rot Disease of Coconut

M. Surulirajan<sup>a++\*</sup>, R. Arunkumar<sup>a#</sup>, V. G. Mathirajan<sup>b†</sup>,  
R. Babu<sup>a‡</sup>, Vinayak Hegde<sup>c^</sup> and Ravi Bhat<sup>c##</sup>

<sup>a</sup> Coconut Research Station, Veppankulam, Tamil Nadu, India.

<sup>b</sup> Agricultural College and Research Institute, Echankottai, Thanjavur, Tamil Nadu, India.

<sup>c</sup> ICAR-Central Plantation Crops Research Institute, Kasaragod, Kerala, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i143052

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101105>

Short Research Article

Received: 01/04/2023  
Accepted: 02/06/2023  
Published: 06/06/2023

## ABSTRACT

Basal stem rot disease is the most challenging disease in coconut crop, caused by *Ganoderma lucidum*. Combating the disease with new generation fungicides is a viable strategy for the promising disease control. Single and combination of new systemic fungicides in different commercial formulations were tested against the *Ganoderma lucidum*. Under *in vitro* study at 100, 250 and 500 ppm concentrations. The results revealed that Hexaconazole 4% + Carbendazim 16% SC, Hexaconazole 5% + Validamycin 2.5%SC and Azoxystrobin 11% + Tebuconazole 18.3% SC

<sup>++</sup> Assistant Professor (Plant Pathology);

<sup>#</sup> Associate Professor (Horticulture);

<sup>†</sup> Associate Professor (Agr. Entomology);

<sup>‡</sup> Professor and Head;

<sup>^</sup> Principal Scientist and Head (Crop Protection);

<sup>##</sup> Programme Coordinator Incharge;

\*Corresponding author: E-mail: [suruliplantpath@gmail.com](mailto:suruliplantpath@gmail.com), [surulirajan.m@tnau.ac.in](mailto:surulirajan.m@tnau.ac.in);

W/W were found superior in inhibiting the mycelial growth of *Ganoderma* as compared to other fungicides. Per cent inhibition indicated the effectiveness of potent fungicides against the pathogen even at lower concentration.

**Keywords:** Basal stem rot; coconut; *Ganoderma lucidum*; fungicide; hexaconazole and in-vitro.

## 1. INTRODUCTION

“Coconut (*Cocos nucifera* Linn.) is the most important successful tropical plantation crop and it provides food, oil, health drink, medicine, fiber, timber, fuel, and varieties of products of commercial importance. Indonesia, Philippines, India and Sri Lanka are the major coconut growing countries. The cultivation of coconut in India during 2016-17 was 2.09 million hectares area with the production of 15339.10 million nuts” (CDB, 2016). “The state Tamil Nadu occupies an area of 0.44 million hectares with production of 4097.23 million nuts” (Saxena, 2017).

“Coconut is affected by various fungal diseases viz., bud rot (*Phytophthora palmivora*), Thanjore Wilt (or) basal stem end rot (*Ganoderma lucidum*), grey blight (*Pestalotiopsis palmarum*) and stem bleeding disease (*Thievoloopsis paradoxa*) occurring in Tamil Nadu. Among them, basal stem rot disease caused by *Ganoderma lucidum* and *G.applanatum* is the most serious disease limiting the coconut production and productivity predominantly occurs in traditional coconut growing areas in East Coastal region, also known as Thanjavur wilt recorded the incidence up to 31 per cent” [1]. Naik [2] mentioned that this disease is also known as *Ganoderma* wilt (Andhra Pradesh) or Tanjavur wilt (Tamil Nadu) or Bole rot or anabe roga (Karnataka) in different parts of India.

“The vertical spread of stem bleeding seen as exudation of reddish-brown viscous fluid is the first sign of the disease. Drooping, drying and falling of leaves, excessive root rot and death of palms are the characteristic symptoms of the basal stem rot. The severely infected palm becomes wilt and unproductive with the formation of sporophore at the base of the palm” [3]. This disease spreads quickly in the ill-maintained coconut gardens.

Snehalatharani et al. [4] assessed “the incidence and spread of coconut basal stem rot disease in different districts of Andhra Pradesh, Karnataka and Tamil Nadu during 2010-15. It was confirmed that Thanjavur district recorded maximum mean per cent incidence of BSR (6.5) followed by Nagapattinam and Thiruvarur in Tamil Nadu”.

Although, many investigations on the field application of disease management practices were tried against this disease, the use of systemic chemicals is highly promising [2], (Karthikeyan et al. 2005), [5-7]. Recently, the aggravation of the disease after the cyclone – Gaja, severe yield reduction and the quest for new systemic chemicals were viewed seriously by the state agriculture research and extension system and coconut growers. This necessitates the *in vitro* evaluation of fungicides to find out the effective new fungicide against *Ganoderma lucidum* for field application. The result of the present work will provide a base to manage basal stem rot of coconut effectively.

## 2. MATERIALS AND METHODS

The present study was done in Coconut Research Station, Veppankulam in 2020. The commercial formulation of thirteen fungicides viz., Famoxadone 16.6% + Cymoxanil 22.1% SC, Cyazafamid 34.5% SC, Mancozeb 35% SC, Kitazin 48% EC, Hexaconazole 4% + Carbendazim 16% SC, Azoxystrobin 11% + Tebuconazole-18.3% SC W/W, Thifluzamide 24% SC, Carbendazim 46.27% SC, Isoprothiolone 40% EC, Difenconazole 25% EC, Azoxystrobin 18.2% w/w + difenoconazole 11.4% w/w SC, Hexaconazole 5% + validamycin 2.5% SC, Hexaconazole 5% + validamycin 2.5% SC and Pencycuron 23.9 % SC were used for the study.

The poisoned food technique [8] was followed to test the efficacy of different fungicides under *in vitro* conditions. The inhibitory effect of thirteen fungicides at 100, 250 and 500 ppm concentrations on the growth of virulent isolate of *Ganoderma lucidum*, (VPM) causing basal stem rot disease on coconut under *in vitro* condition was evaluated by poisoned food technique. Each chemical was replicated four times with proper control.

“The required quantities of fungicides were added into the sterilized Potato Dextrose Agar (PDA) medium to give required concentration and poured separately into each sterilized Petri plates under aseptic conditions. The Petri plates were inoculated with 8 mm mycelia disc from

seven days old culture of the fungus and incubated at  $28 \pm 2^\circ\text{C}$ . Simultaneously, a control was maintained without adding fungicide by growing the fungus only on PDA medium. Three replications were kept for each treatment. The observations were made on the diameter of mycelial growth of the fungus. The per cent inhibition in mycelial growth was calculated by using the following formula: Each treatment was replicated four times in a completely randomized design. The data obtained were statistically analyzed [9] and the treatment means were compared by Duncan's Multiple Range Test (DMRT)".

Per cent inhibition over control =  $\frac{DC - DT}{DC} \times 100$  CWhere,

PI = Per cent inhibition

DC = Mean diameter (cm) of fungal growth in control DT = Mean diameter (cm) of fungal growth in treatment.

### 3. RESULTS AND DISCUSSION

The inhibitory effect of thirteen fungicides viz., Famoxadone 16.6%+ Cymoxanil 22.1% SC, Cyazafamid 34.5% SC, Mancozeb 35% SC, Kitazin 48% EC, Hexaconazole 4% + Carbendazim 16% SC, Azoxystrobin 11% + Tebuconazole-18.3% SC W/W, Thifluzamide 24% SC, Carbendazim 46.27% SC, Isoprothiolone 40% EC, Difenconazole 25% EC, Azoxystrobin 18.2% w/w + difenoconazole 11.4% w/w SC, Hexaconazole 5% + validamycin 2.5% SC, Hexaconazole 5% + validamycin 2.5% SC and Pencycuron 23.9 % SC tested against *Ganoderma lucidum* causing basal stem rot disease on coconut under *in vitro* by poisoned food technique at 100, 250 and 500 ppm concentrations and the growth of *Ganoderma lucidum*, was evaluated.

The results revealed that among thirteen fungicides tested, Hexaconazole 4% + Carbendazim 16% SC, Azoxystrobin 11% + Tebuconazole-18.3% SC W/W and Hexaconazole 5% + validamycin 2.5% SC were found superior recording 100 per cent inhibition of *Ganoderma lucidum* at all the concentrations tested and showed superior over the other fungicides tested under *in vitro*.

Anbalagan and Shanmugam [10] evaluated the fungicide Tridemorph at 500 ppm against *G. lucidum in vitro* and was found effective in inhibiting the spread of the fungus. Srinivasulu et al. [11] conducted *in vitro* screening of the fungicides viz., Bordeaux mixture, Tridemorph, Bitertenol, Copper oxychloride and Hexaconazole against *G. lucidum* and were found effective in inhibiting the growth of *G. lucidum* and *G. applanatum* and also found inhibitory to *Trichoderma viride*.

Baloch et al. [12] reported that "the fungicide tebuconazole + trifloxystrobin inhibited the mycelial growth of *L. theobromae* mycelium at 100 ppm. showed structural abnormalities in response to fungicide treatment which resulted in shriveling. Ushamalini et al. [13] also screened fifteen fungicides for their inhibitory effect on the mycelial growth of *L. theobromae* at different concentrations and reported that systemic fungicide carbendazim, recorded 100 percent inhibition over control even at lower concentration of 50 ppm". The present study of *In-vitro* experiment evaluated the fungicide chemicals in new combination against *Ganoderma lucidum* and all the treatments are found significant at 5% in three concentrations viz., 100ppm (F-Probability is 3.5 & Co-efficient of Variation is 9.146), 250ppm (F-Probability is 1.04 & Co-efficient of Variation is 0.706) and 500ppm (F-Probability 3.26 & Co-efficient of Variation is 0.852). Hundred per cent mycelial inhibition was achieved by three new combinations fungicides viz., Hexaconazole 4% + Carbendazim 16% SC, Azoxystrobin 11%+ Tebuconazole-18.3% SC W/W and Hexaconazole 5% + validamycin 2.5% SC (Table 1). Similar results were recently reported by Thangeshwari et al. (2019) with the screening of twelve fungicides and new fungicide combinations against *Ganoderma lucidum* and found that Tebuconazole 25.9 per cent EC, Tetraconazole 3.8 per cent EW, Tebuconazole + Trifloxystrobin 50 per cent + 25 per cent WG, Hexaconazole 5 per cent EC, Difenconazole 25 per cent EC, Thiram + Carboxin 37.5 + 37.5 WS and Propiconazole 25 EC have recorded 100 per cent inhibition of *Ganoderma lucidum* and showed superiority over other tested fungicides under *in vitro*. Karunanithi et al. [5] also conducted *in vitro* studies and recorded the suppression of the fungus by 29 botanicals.

**Table 1. In-vitro evaluation of combination systemic fungicides against *Ganoderma lucidum***

Treatments	Fungicides	Mycelial growth of <i>Ganodermaspp.</i> 9 days after inoculation (in mm)			% Inhibition over control
		100ppm	250ppm	500ppm	
T <sub>1</sub>	Famoxadone 16.6% + Cymoxanil 22.1% SC	90.00 <sup>a</sup> (71.67)	90.00 <sup>a</sup> (71.67)	89.68 <sup>a</sup> (71.26)	0.361
T <sub>2</sub>	Cyazafamid 34.5% SC	90.00 <sup>a</sup> (71.67)	90.00 <sup>a</sup> (71.67)	90.00 <sup>a</sup> (71.67)	00.00
T <sub>3</sub>	Mancozeb 35% SC	68.75 <sup>b</sup> (56.01)	68.13 <sup>b</sup> (55.63)	66.38 <sup>b</sup> (54.56)	26.25
T <sub>4</sub>	Kitazin 48% EC	15.38 <sup>f</sup> (23.09)	11.20 <sup>f</sup> (19.55)	10.39 <sup>f</sup> (18.80)	88.46
T <sub>5</sub>	Hexaconazole 4% + Carbendazim 16% SC	00.00 <sup>h</sup> (00.00)	00.00 <sup>i</sup> (00.00)	00.00 <sup>i</sup> (00.00)	100.00
T <sub>6</sub>	Azoxystrobin 11% + Tebuconazole-18.3% SC W/W	1.50 <sup>gh</sup> (7.03)	00.00 <sup>i</sup> (0.00)	00.00 <sup>i</sup> (0.00)	100.00
T <sub>7</sub>	Thifluzamide 24% SC	18.06 <sup>f</sup> (25.15)	10.31 <sup>g</sup> (18.73)	12.31 <sup>e</sup> (20.54)	86.32
T <sub>8</sub>	Carbendazim 46.27% SC	6.13 <sup>g</sup> (14.33)	4.13 <sup>h</sup> (11.73)	0.81 <sup>h</sup> (5.16)	99.10
T <sub>9</sub>	Isoprothiolone 40% EC	35.13 <sup>d</sup> (36.35)	23.43 <sup>d</sup> (28.95)	15.19 <sup>d</sup> (22.94)	83.13
T <sub>10</sub>	Difenoconazole 25% EC	56.44 <sup>c</sup> (48.70)	46.49 <sup>c</sup> (42.99)	35.50 <sup>c</sup> (36.57)	60.56
T <sub>11</sub>	Azoxystrobin 18.2% w/w + difenoconazole 11.4% w/w SC	25.19 <sup>e</sup> (30.13)	21.00 <sup>e</sup> (27.27)	3.78 <sup>g</sup> (11.21)	95.81
T <sub>12</sub>	Hexaconazole 5% + validamycin 2.5% SC	00.00 <sup>h</sup> (00.00)	00.00 <sup>i</sup> (00.00)	00.00 <sup>i</sup> (00.00)	100.00
T <sub>13</sub>	Pencycuron 23.9 % SC	90.00 <sup>a</sup> (71.67)	90.00 <sup>a</sup> (71.67)	90.00 <sup>a</sup> (71.67)	00.00
T <sub>14</sub>	Control	90.00 <sup>a</sup> (71.67)	90.00 <sup>a</sup> (71.67)	90.00 <sup>a</sup> (71.67)	00.00
	<b>CD@5%</b>	<b>5.465</b>	<b>0.391</b>	<b>0.437</b>	
	<b>CV</b>	<b>9.146</b>	<b>0.706</b>	<b>0.852</b>	

Figures in parenthesis are transformed values

#### 4. CONCLUSION

In-vitro experiment evaluated the fungicide chemicals in new combination against *Ganoderma lucidum*. The hundred per cent mycelial inhibition of *Ganoderma lucidum* was achieved by three new fungicide combinations viz., hexaconazole 4% + carbendazim 16% sc, azoxystrobin 11% + tebuconazole-18.3% sc w/w and hexaconazole 5% + validamycin 2.5%sc.

#### ACKNOWLEDGEMENTS

ICAR-AICRP on Palms is thankfully acknowledged by the authors for financial assistance

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Bhaskaran R, Ramanathan T. Occurrence and spread of Thanjavur wilt disease of coconut. Indian Coconut Journal. 1984; 15(6):12–14.
2. Naik RG. Chemical control of basal stem rot of coconut. Agricultural Science Digest 2001;21(4):249.
3. Bhaskaran R, Rethinam P, Nambiar KKN. Thanjavur wilt of coconut. Journal of Plantation Crops. 1989;17(2):69–79.
4. Snehalatharani A, Devappa V, Rajappan K and Maheswarappa HP. Coconut basal stem rot is manageable. Indian Horticulture. 2014a;59(6):37-38.
5. Karunanithi K, Sarala L, Rabindran R, Sabitha Doraisamy, Rajarathinam S, Khan HH. Effect of plant products on the management of basal stem rot (*Ganoderma*) of coconut. Indian Coconut Journal. 2007;38(2):13–5.

6. Surulirajan M, Rajappan K, Kumar NS, Annadurai K, Kumar KJ, Asokhan M. Management of basal stem rot disease in coconut through bio-inoculants and chemicals. *Indian Journal of Tropical Agriculture*. 2014;32(3-4):407-14.
7. Snehalatharani A, Chalapathi Rao NBV, Ramanandam G, Maheswarappa HP, Jose CT and Padma E. Bio-control based integrated disease management of basal stem rot disease of coconut. *Journal of Plantation Crops*. 2016;44(1): 62-66.
8. Schmitz H. Food Poisoned Technique. *Industrial Engineering Chemical Analyst Education*. 1930;361-363.
9. Gomez KA, Gomez AA. *Statistical procedures for agricultural research*: John Wiley & Sons. 2<sup>nd</sup> Edition, New York; 1984.
10. Anbalagan R, Shanmugam N. Studies on *Ganoderma lucidum* (Leys) Karst. associated with Thanjavur wilt of coconut. In: *Proceedings of All India Thanjavur wilt of coconut*. In: *Proceedings of All India Nadu*. 1984;24-29.
11. Srinivasulu B, Aruna K, Krishna Prasadji J, Rajamannar, Sabitha Doraisamy, Rao DVR, Hameed Khan H. Prevalence of basal stem rot disease of coconut in coastal agro-ecosystem of AP. *Indian Coconut Journal*. 2002;XXXIII:23-26.
12. Baloch A, Abro MA, Jatoi GH, Abro NA, Dad S, Rafique M, Ali R, Memon S, Mangi N. *In-vitro* evaluation of different fungicides and the bio-control agent *Trichoderma harzianum* against *Botryodiplodia theobromae*, the cause of guava decline. *Pakistan J. Biotechnol*. 2017;14:491-496.
13. Ushamalini C, Ashok kumar P, Parthasarathy S. Efficacy of fungicides for the management of coconut leaf blight. *Pl. Dis. Res*. 2019;34(1):29-35.

© 2023 Surulirajan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/101105>