



Effect of Organic Manures and Inorganic Fertilizers on Growth, Yield and Quality of Onion (*Allium cepa*) var. Nashik Red

Kalle Preethi^{a*}, Anita Kerketta^{b#}, Vijay Bahadur^{b‡} and Sanam Rajkumar^{a*}

^a Department of Horticulture (Vegetable Science), NAI, SHUATS, Prayagraj-211007, (U.P.), India.

^b Department of Horticulture, NAI, SHUATS, Prayagraj-211007, (U.P.), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i1131130

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/90627>

Original Research Article

Received 04 June 2022

Accepted 08 August 2022

Published 12 August 2022

ABSTRACT

An experiment was conducted to find out the effect of organic manures and inorganic fertilizers on growth, yield and quality of onion (*Allium cepa* L.) var. Nashik Red during 2021-2022 in Rabi season (Nov-Apr) at Horticulture Research Farm, Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Allahabad (Uttar Pradesh). Total 13 treatments were replicated thrice in Randomized Block Design (RBD). The results showed that treatment T4 (75% NPK + 25% Poultry manure) was recorded maximum in growth and quality parameters i.e., plant height (62.29 cm), maximum number of leaves (10.38), maximum TSS (14.58 °Brix) and maximum ascorbic content of (12.49). whereas, the yield parameters were maximum in treatment T8 (50% NPK + 50% Poultry manure) was recorded with maximum weight of fresh bulb weight (181.70 g), maximum weight of cured bulb (168.07 g), maximum bulb diameter (8.63 cm), maximum yield per plot (23.09 kg) and maximum yield per hectare (15.36 t).

Keywords: Recommended dose of fertilizers (RDF); nitrogen- phosphorus- potassium (NPK); farm yard manure (FYM); total soluble solids (TSS).

*M.Sc. Scholar;

Assistant Professor;

‡ Associate Professor;

*Corresponding author: E-mail: preethij1707@gmail.com;

1. INTRODUCTION

Onion (*Allium cepa*) belongs to the family Amaryllidaceae and it is known as "Pyaj" in Hindi. It occupies very important position among spices crops grown all over the world. Onion ranks first position in area under spices in the country and occupies an area of 0.39 million hectare with the second rank in production i.e., 4.58 million tonnes. In the world production Indian shares 12% after China of the world vegetable production. Onion is cultivated throughout India but Maharashtra, Gujrat, Karnataka, Uttar Pradesh, Andhra Pradesh, Tamil Nādu, M.P., Bihar, and Assam are the important onion producing states. Now-a-days consumer's preferences have also shifted away from cereals and moved towards vegetables for balanced diet [1]. Among the bulb crops onion is the most important bulb vegetable crop and an essential commodity. Onion is considered as cool season crop and grows normally in winter seasonal most all over India. Onion has great value as a medicinal plant. Pungency in onion is due to volatile oil allylprophyl disulphide ($C_6H_{12}O_2$), which acts as gastric stimulant and promote digestion. Onions are highly prized for their medical characteristics, which include antiseptic and antibacterial capabilities. It relieves heat and is used to treat hysterical outbursts and faintness. The study was undertaken to estimate the effect of organic manures and inorganic fertilizers on growth, yield and quality of onion (*Allium cepa*). The organic manures help in the improvement of soil physical and physiological structure whereas the inorganic fertilizers can improve yield of the crop

and the total nutrient availability. Therefore, application of organic and inorganic in different combinations was carried out in aim to assess the suitable combination to maintain both soil health and yield of crop as well.

2. MATERIALS AND METHODS

The experiment was carried out *Rabi* season 2021 at Research farm in Department of Horticulture, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P). The experiment was laid out in Randomized Block Design (RBD) with thirteen treatments, those were replicated thrice, Nashik red variety of onion was selected for the study. Which was planted during *Rabi* season in 2021 at 15cm X 10cm spacing and organic manures and inorganic fertilizers were applied in regular intervals. The treatments details were mentioned in table 1. The observations on growth parameters i.e., plant height (cm), number of leaves, bolting percent were recorded from randomly selected plants from each and every plot at various vegetative growth stages. Yield parameters i.e., days to maturity, weight of fresh bulb (g), weight of cured bulb (g), bulb diameter (cm), number of bulbs per kg, yield per plot (kg) and yield per hectare (ton) were also recorded. Quality parameters i.e., Total soluble solids ($^{\circ}$ Brix) and ascorbic acid content were recorded. The data was analyzed using the ANOVA approach and a significant difference between the treatments means was checked against the critical difference at a 5% level of significance.

Table 1. Treatment details

Treatments symbols	Treatment combinations
T ₀	100%RDF (100:50:50 kg NPK)
T ₁	25%NPK+75% FYM
T ₂	25%NPK+75% Vermicompost
T ₃	25%NPK+75% goat manure
T ₄	25% NPK+75% poultry manure
T ₅	50% NPK+50% FYM
T ₆	50% NPK+50% vermicompost
T ₇	50% NPK+50% goat manures
T ₈	50% NPK+50% poultry manure
T ₉	75% NPK+25% FYM
T ₁₀	75% NPK+25% vermicompost
T ₁₁	75% NPK+25% goat manures
T ₁₂	75%NPK + 25% poultry manure

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

From the below Table 2, the results revealed that significantly the maximum plant height (62.29 cm), was recorded in the treatment T4 (75% NPK + 25% Poultry manure) followed by T6 with (60.85 cm), while the minimum plant height was recorded in T0 *i.e.*, 100% RDF (100:50:50 NPK) (57.33 cm). The increase in plant height is due to the integrated use of nutrient helpful in development of rapid cell division and cell elongation in meristematic region of plant due to production of plant growth substance and this may be due to abundant supply of plant nutrients and nitrogen which led in the growth of onion. In general, height of the plant was found to be more when plots applied with 75 per cent RDF (inorganic source) + 25 per cent organic manures. The results obtained in the present study are supported by the findings reported by Kannan [2], indicated that combination of 12.5 t/ha poultry manure with 50 kg N/ha increased plant height.

The maximum number of leaves were recorded in treatment T4 (75%NPK + 25% Poultry manure (10.38) statistically at par with T2 (75%NPK + 25%Vermicompost) (10.12) followed by 50% NPK + 50% Farmyard manure (9.93) while the minimum number of leaves were recorded in 100% RDF (50%NPK + 50%Goat manure) (8.44). It can be seen that organic source of nutrient along with inorganic fertilizer showed

better response in terms of increasing number of leaves per plant as compared to inorganic source alone. Significant differences were observed at all the treatments at all growth stages of observation. More number of leaves per plant was obtained when the plants applied with organic manures along with inorganic fertilizer as compared to application of chemical fertilizer alone. This effect could be attributed to the quick fermentation and solubilization effect of plant nutrients of poultry manure. Also, poultry manure might have enhanced use efficiency of chemical fertilizer. The result obtained in the present study are supported by the findings reported by Khalil et al., [3] observed that chicken manure and inorganic fertilizers were more effective than FYM in increasing leaf number per plant in onion. Jaythilake et al., [4] reported that integrated effect of inorganic fertilizers, organic manure and biofertilizers gave a greater number of leaves in onion.

The minimum bolting was recorded in (T8) 50%NPK + 50% Poultry manure (5.86 %) followed by (T5) 50%NPK + 50%Farm yard manure (6.09%) while the maximum bolting was recorded in (T0) 100%RDF (100:50:50 NPK) (10.61%). Bolting in onion crop is most undesirable character which adversely affects the quality and storage life of onion bulbs. Bolting is usually variable from year to year and among cultivars. These results were in accordance with those of Peterson et al., [5] stated that premature seed stalk development decreased with addition of nitrogen. Diaz-Perez et al., [6] reported that

Table 2. Effect of organic manures and inorganic fertilizers on growth parameters of onion

Treatments symbols	Treatment combinations	Plant height (cm)	Number of leaves per plant	Bolting (%)
T ₀	100% RDF	57.33	9.05	10.61
T ₁	75%NPK + 25% Farm yard manure	57.09	9.43	6.65
T ₂	75%NPK + 25% Vermicompost	60.85	10.12	6.16
T ₃	75%NPK + 25% Goat manure	60.49	8.80	8.33
T ₄	75%NPK + 25% Poultry manure	62.29	10.38	7.09
T ₅	50%NPK + 50% Farm yard manure	60.12	9.93	6.09
T ₆	50%NPK + 50% Vermicompost	60.15	9.22	6.12
T ₇	50%NPK + 50% Goat manure	59.76	8.44	8.14
T ₈	50%NPK + 50% Poultry manure	59.52	8.65	5.86
T ₉	25%NPK + 75%Farm yard manure	56.97	9.09	7.60
T ₁₀	25%NPK + 75% Vermicompost	57.84	8.61	7.91
T ₁₁	25%NPK + 75% Goat manure	57.09	8.52	8.86
T12	25%NPK + 75% Poultry manure	58.56	8.84	8.75
	F-Test	S	S	S
	C.D at 5%	0.61	0.29	0.13
	S.Ed.	0.30	0.14	0.06
	C.V.	0.61	1.88	1.05

the N fertilization rates increased bolting N fertilization rates applied ranged from the infra-optimal to the super optimal from (102 to 302/ha N).

3.2 Yield Parameters

The results revealed that, significantly minimum number of days for maturity (106.30) were recorded in treatment T8 (50%NPK + 50%Poultry manure) which is followed by T5 (50%NPK + 50%Farm yard manure) (108.12) whereas, maximum days for maturity were recorded in T0 with 100%RDF (100:50:50 NPK) 117.82 days. The probable reason for early maturity may be that the hormones and organic acid secreted by organic manures during decomposition which enhanced early maturity. The results in present study are supported by the findings of Kumaran [7]. Raut [8] and Barekar [9] in onion.

The maximum weight of fresh bulb was recorded in T8 (50% NPK + 50% Poultry manure) 181.70 g which is followed by T2 (75% NPK + 25% Vermicompost) 171.27 g whereas, the minimum weight of fresh bulb was recorded in T0 100%RDF (100:50:50 NPK) with 100.25 g.

The maximum weight of cured bulb was recorded in T8 (50% NPK + 50% Poultry manure) 168.07 g which is followed by T2 (75% NPK + 25% Vermicompost) 159.59 g whereas, the minimum weight of cured bulb was recorded in T0 100% RDF (100:50:50 NPK) 96.13 g. The treatments receiving 50% RDF + 50% poultry manure produced more average bulb weight. The increase in average bulb weight was mainly due to increased bulb diameter, the relative significant increase in the number of green leaves due to the combined effects in those treatments may have helped to accumulate more carbohydrates resulting in increased diameter of the bulb which results in increase in fruit weight. Similar results were obtained by Singh [10] and Lal et al., [11].

The maximum bulb diameter was recorded in T8 (50%NPK + 50%Poultry manure) with 8.63 cm which is followed by T5 (50% NPK + 50% Farm yard manure) with 8.27 cm whereas, the minimum bulb diameter was recorded in T0 100% RDF (100:50:50 NPK) 6.43 cm. The total number of leaves the plants generated had a direct impact on the size of the bulb. Given that the bulb is the onion's storage organ and that whatever carbohydrates were produced in the

leaves were stored in the bulbs, resulting in larger sized bulbs as indicated, the relative significant increase in the number of green leaves due to the combined effects in those treatments may have helped to accumulate more carbohydrates resulting in increased diameter of the bulb. These results were in agreement with those reported by Nagaraju et al., [12], and Mondal et al., [13] in onion.

The minimum number of bulbs per kg were recorded in T8 (50% NPK + 50% Poultry manure) 5.91, followed by treatment T10 (25% NPK + 75% Vermicompost) 7.48 whereas, the maximum number of bulbs per kg were recorded in T2 (75% NPK + 25% Vermicompost) with 10.34. This might be because of more fresh and cured weight of bulbs accommodating less number of bulbs per kg.

The maximum yield per plot were recorded in T8 (50% NPK + 50% Poultry manure) with 23.09 kg which is followed by T5 (50% NPK + 50% Farm yard manure) with 21.41 kg whereas, the minimum yield per plot were recorded in T0 100% RDF (100:50:50 NPK) with 14.24 kg. The maximum yield per hectare were recorded in T8 (50% NPK + 50% Poultry manure) with 15.39 ton which is followed by T5 (50% NPK + 50% Farm yard manure) with 14.27 ton whereas, the minimum yield per hectare were recorded in T0 100% RDF (100:50:50 NPK) with 9.49 ton. According to the highest plants and leaf count, the increase in yield appeared to be the result of higher plant growth.

3.3 Quality Parameters

The maximum TSS were recorded in T4 (75%NPK + 25% Poultry manure) with 14.58 which is statistically at par with T8 (50% NPK + 50% Poultry manure) with 14.21 followed by T5 (50% NPK + 50% Farm yard manure) with 14.10 whereas, the minimum TSS were recorded in T0 100% RDF (100:50:50 NPK) with 12.89. The total soluble solids were observed to increase over the prescribed dose of inorganic fertilizers in all other treatment combinations of organic and inorganic fertilizers with respective levels.

The maximum ascorbic acid content was recorded in T4 (75% NPK + 25% Poultry manure) with 12.49 which is followed by T5 (50% RDF + 50% Farm yard manure) with 11.76 whereas, the minimum ascorbic acid content was recorded in T3 (75% NPK + 25% Goat manure) with 10.07. This might be due to physiological

Table 3. Effect of organic manures and inorganic fertilizers on yield parameters of onion (*Allium cepa*)

Treatments symbols	Treatment combinations	Days to Maturity	Weight of fresh bulb (g)	Weight of cured bulb (g)	Bulb diameter(cm)	Number of bulbs per kg	Yield per plot (kg)	Yield per hectare (ton)
T ₀	100% RDF	117.82	100.25	96.13	6.43	7.35	14.24	9.49
T ₁	75%NPK + 25% Farm yard manure	110.13	129.07	120.15	7.90	6.50	19.76	13.17
T ₂	75%NPK + 25% Vermicompost	112.61	171.27	159.59	7.92	10.34	18.41	12.27
T ₃	75%NPK + 25% Goat manure	111.55	137.09	112.15	7.68	6.58	18.59	12.39
T ₄	75%NPK + 25% Poultry manure	109.37	144.58	134.30	8.07	6.31	19.45	12.96
T ₅	50%NPK + 50% Farm yard manure	108.12	151.57	147.73	8.27	6.41	21.41	14.27
T ₆	50%NPK + 50% Vermicompost	113.20	146.54	141.86	7.67	6.92	18.40	12.26
T ₇	50%NPK + 50% Goat manure	113.46	142.82	138.17	7.69	6.53	17.31	11.53
T ₈	50%NPK + 50% Poultry manure	106.30	181.70	168.07	8.53	5.91	23.09	15.39
T ₉	25%NPK + 75%Farm yard manure	111.70	140.38	135.71	6.87	7.38	19.16	12.77
T ₁₀	25%NPK + 75% Vermicompost	111.62	137.50	133.41	7.47	7.48	17.20	11.46
T ₁₁	25%NPK + 75% Goat manure	112.54	140.78	135.64	7.39	7.20	16.92	11.27
T ₁₂	25%NPK + 75% Poultry manure	110.36	149.11	144.99	7.70	6.89	17.58	11.71
	F-Test	S	S	S	S	S	S	S
	C.D at 5%	0.36	2.54	0.65	0.10	0.09	0.25	0.17
	S.Ed.	0.18	1.23	0.32	0.05	0.05	0.12	0.08
	C.V.	0.19	1.05	0.28	0.75	0.79	0.81	0.80

Table 4. Effect of organic manures and inorganic fertilizers on quality parameters in onion

Treatments symbols	Treatment combinations	TSS	Ascorbic acid
T ₀	100% RDF	12.89	10.09
T ₁	75%NPK + 25% Farm yard manure	13.48	11.30
T ₂	75%NPK + 25% Vermicompost	13.15	11.20
T ₃	75%NPK + 25% Goat manure	13.15	10.07
T ₄	75%NPK + 25% Poultry manure	14.58	12.49
T ₅	50%NPK + 50% Farm yard manure	14.10	11.76
T ₆	50%NPK + 50% Vermicompost	13.16	10.29
T ₇	50%NPK + 50% Goat manure	12.97	10.26
T ₈	50%NPK + 50% Poultry manure	14.21	11.59
T ₉	25%NPK + 75%Farm yard manure	13.82	11.40
T ₁₀	25%NPK + 75% Vermicompost	13.12	10.11
T ₁₁	25%NPK + 75% Goat manure	12.92	10.09
T ₁₂	25%NPK + 75% Poultry manure	13.50	11.01
	F-Test	S	S
	C.D at 5%	0.41	0.09
	S. Ed.	0.20	0.04

influence of poultry manure and FYM in combination with inorganic sources of nutrients on activity of number of enzymes and due to more energy and food material available to the bulbs due to strong vegetative growth. Similar results were also reported by Mondal et al., [13] in onion, Kumaran et al., [7] in tomato [14].

4. CONCLUSION

Among the various combinations, it was concluded that overall performance of the treatment T₄ which received 75 per cent NPK through chemical fertilizers along with 25 per cent poultry manure per hectare as organic manure enhanced vegetative growth and quality of onion.

These results are based on one season study and for confirmation few more experimental trials are needed.

ACKNOWLEDGEMENT

The work was supported by Advisor Dr. Anita Kerketta and all faculty members, Department of Horticulture, Naini Agricultural institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, (U.P.) India for providing field and assistance in conducting the research trail.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Mittal S. Can horticulture be a success story for India? (No. 197). Working Paper; 2007.
- Kannan B. Effect of different source of nitrogen on nutrient uptake, quality and yield of Bhendi (*Abelmoschus esculentus* L. Moench). M. Sc (Ag.), Tamil Nadu Agricultural University, Coimbatore; 1990.
- Khalil MY, Naguib NY, ElSherbeny SE. Effect of (*Tagetes erecta* L.) to some foliar application under compost levels. Arab University. Journal of Agricultural Science. Ain Shams University, Cairo. 2002; 10(3):939-964.
- Jayathilake PKS, Reddy IP, Srihari D, Neeraja G, Ravinder Reddy. Effect of nutrient management on growth, yield and yield attributes of rabi onion (*Allium cepa* L.) Vegetable Science. 2002;29(2):184-185.
- Paterson DR, Black Hurst HT, Siddique HS. Some effects of nitrogen and phosphoric acid on premature seed stalk development, yield and composition of three onion varieties. Proc. American Society of Horticultural Science. 1960; 76:460-467.
- Díaz-Pérez JC, Purvis AC, Paulk JT. Bolting, yield, and bulb decay of sweet onion as affected by nitrogen fertilization. Journal of the American Society for Horticultural Science. 2003; 128(1):144-149.

7. Kumaran S, Natarajan S, Thamburaj S. Effect of organic and inorganic fertilizer on growth yield and quality of tomato. South Indian Horticulture. 1998;46(3-4):203-205.
8. Raut SP. To study the availability of export quality pods of okra as influenced by integrated nutrient management and spacing. M.Sc. (Agri.) Thesis, MAU, Parbhani; 1998.
9. Barekar NN. Effect of integrated nutrient management on growth and yield of chilli (*Capsicum annum* L.) M.Sc. (Agri.) Thesis, Dr. P.D.K.V., Akola, (M.S.); 2000.
10. Singh L, Bhonde BR, Mishra UK. Effect of different organic manures and inorganic fertilizers on yield and quality of rabi onion. Newsletter, N.H.R.D.F. 1997;17(3):1-3.
11. Lai S, Yadav AC, Mangal JB, Avtarsingh, Batra VK. Effect of FYM and irrigation level on growth and yield of onion cv. Hissar 2. Haryana Journal of Horticultural Science. 2002;31(3-4):256-258.
12. Nagaraju R, Haripriya K, Rajalingam GV, Shriramachandrasekaran V, Mohindeen MK. Effect of YAM on growth and yield of aggregatum onion (*Allium cepa* L. Var. Aggregatum Don). South Indian Horticultural. 2000;48(1/6):40-45.
13. Mondal SS, Debabrata Acharya, Amp Ghosh, Thapa U. Integrated management of organic and inorganic sources of nutrient to improve productivity and qualitative characters of rice and onion in rice-onion cropping sequence. Environmental & Ecology. 2004;22(1):125-128.
14. Rather SA, Ahmed N, Chattoo MA. Respons of onion to microbial inoculation and chemical nitrogen. Haryana Journal of Horticultural Science. 2003;32(3/4):270-271.

© 2022 Preethi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/90627>