



## **Evaluation of Insulin, Malondialdehyde and Blood Pressure in Male Obese Individuals in Nnewi and Subsequent Effect of Green Tea Supplementation**

**Onuora Ifeoma Joy<sup>1\*</sup>, Meludu Samuel Chukwuemeka<sup>2</sup>, Dioka Chudi Emmanuel<sup>1</sup>, Obi-Ezeani Chikaodili Nwando<sup>1</sup> and Usman Saheed Opeyemi<sup>1</sup>**

<sup>1</sup>Department of Chemical Pathology, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria.

<sup>2</sup>Department of Human Biochemistry, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria.

### **Authors' contributions**

*This work was carried out in collaboration between all authors. Authors OIJ and MSC designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors OIJ and OECN managed the analyses of the study. Authors USO and DCE managed the literature searches. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/AJRIMPS/2018/41153

#### Editor(s):

(1) John Yahya I. Elshimali, Professor, Department of Pathology and Oncology, UCLA School of Medicine, Charles R. Drew University of Medicine and Science, California, USA.

#### Reviewers:

(1) Dhastagir sheriff, Benghazi University, Libya.

(2) Mra Aye, Melaka Manipal medical College, Melaka, Malaysia.

Complete Peer review History: <http://www.sciencedomain.org/review-history/24412>

**Original Research Article**

**Received 14<sup>th</sup> February 2018**  
**Accepted 22<sup>nd</sup> April 2018**  
**Published 1<sup>st</sup> May 2018**

### **ABSTRACT**

**Background:** Obesity is a major public health issue worldwide, contributing to increased cardiovascular diseases, diabetes, insulin resistance and oxidative stress. This is due to sedentary lifestyles; poor dieting and low consumption of antioxidant supplement (example green tea). The objective of this study was to evaluate the level of fasting blood sugar, insulin, insulin resistance blood pressure and MDA in obese subjects and subsequent effect of green tea at 6weeks and 12weeks supplementation.

**Methods:** This was a cross sectional and interventional study. In the cross sectional study, 88 obese subjects (46 class I and 42 class II obese) and 50 normal weight subjects (control) were recruited. In the interventional study, 20 male obese subjects were randomly selected and were given 200ml of commercially prepared green tea. Fasting blood samples were collected before the

\*Corresponding author: E-mail: ifeomajoyonuorah@gmail.com, ijonuora@yahoo.com;

intervention (baseline), at 6weeks and 12weeks of intervention and were later analyzed by standard method Enzyme Linked immunoassay and colorimetric method. It was analysed statistically using SPSS version 23.0.

**Results:** There were significant increases in the mean levels of HOMA-IR, systolic and diastolic blood pressures, fasting plasma glucose and insulin in obese subjects (class II and class I obese) when compared with control group ( $P<0.05$ ), likewise in Class II obese when compared with Class I obese ( $P<0.05$ ) while in the case of MDA, there was a significant increase only in Class II obese subjects when compared with the normal weight subjects ( $P<0.05$ ). Green tea supplementation significantly reduced the mean level of MDA, fasting plasma glucose, weight, HOMA-IR and blood pressure at 12weeks of intervention while only Insulin and waist circumference were significantly reduced at 6weeks and 12weeks of intervention.

**Conclusion:** In conclusion, obesity is the major cause of diabetes, high blood pressure and insulin resistance. Green tea could be beneficial to diabetic patients and obese hypertensives. Green tea compounds- phytochemicals could be beneficial as one of the components of their diet.

*Keywords: Obesity; green tea; blood pressure; insulin resistance; oxidative stress.*

## 1. INTRODUCTION

Obesity is a medical condition in which excess body fat accumulates to the extent that it may have a negative effect on health. Recent studies have shown that obesity-associated risk factors depend not on excess body weight *per se*, but rather on the regional distribution of the excess body fat. In light of this, it is now well recognized that abdominal fat is a significant risk factor for obesity-associated diseases; in fact, visceral fat accumulation stimulates pro-oxidant and pro-inflammatory states [1]. In obesity, modulation of metabolic pathways plays critical roles in the pathogenesis of many diseases [2]. There is a strong positive association between obesity and type II diabetes, dyslipidaemia, cardiovascular disease and hypertension [3]. Hence, creation of appropriate strategies to reduce weight, insulin resistance, oxidative stress and to increase total antioxidant capacity in obese, have remain the focus of this study.

Recent studies on humans show that green tea has many health benefits including reduced risk of cardiovascular disease and some cancers, anti-effects on blood pressure, weight loss, antiviral and antibacterial activities, anti-mutagenic, anti-inflammatory and reduce insulin resistance [4]. Green tea contain appreciable amounts of phytochemicals especially catechins that further comprised of different chemical moieties that include epigallocatechin-3-gallate (EGCG), epicatechin (EC), epicatechin-3-gallate (ECG), epigallocatechin (EGC). Amongst these, EGCG is present in higher amounts and considered to be an effective antioxidant. In the recent era, diet based therapy has been revitalized globally and people are adopting the

approach of using natural products as an intervention against various ailments [5]. Keeping in view the health challenges associated with obesity, limited research and controversial findings on effect of green tea on ameliorating blood pressures, oxidative stress and diabetes, this study aims at evaluating these parameters in obesity and subsequent effect of green tea supplementation.

## 2. METHODS

### 2.1 Study Design

The research was carried out at Nnewi, Anambra state, Nigeria and biochemical analyses were performed at Nnamdi Azikiwe Teaching Hospital (NAUTH) Nnewi, Anambra State, Eastern Nigeria. This hospital was chosen because they have competent personnel (Medical laboratory scientist) and equipment. In the cross sectional study, 88 obese subjects (46 class I and 42 class II obese) and 50 normal weight subjects (control) were recruited. In interventional study, 20 male obese subjects were randomly selected and were given green tea.

### 2.2 Source of Green Tea

The green tea was obtained from Lipton Company (Unilever Ghana Ltd (GH) and was of the same brand and batch number 16252 with NAFDAC Reg. NO: B1-8866. Phytochemical analysis on the green tea was performed on a BUCK M910 gas chromatography equipped with a flame ionization detector according to Kelly D; Nelson R; [6] to note the concentration of active ingredients (phenol) present.

### 2.3 Preparation of Green Tea

Two (2) green tea bags, each weighing 1.6 g were dissolved in 200 ml of boiled water and this was left to dissolve for 5mins before consumption. The green tea was taken once daily for 12weeks (3 months).

### 2.4 Inclusion Criteria and Exclusion Criteria

Subjects recruited were between the ages of 29 and 47years with body mass index of 30 - 35 Kg/m<sup>2</sup> (for Class I obese), 35 – 40 kg/m<sup>2</sup> (class II obese) and 19-24.9 Kg/m<sup>2</sup> for non-obese (controls). Apparently healthy individuals who were not on any medications for diabetes, hypertension and other CVD were recruited. Subjects on alcohol, cigarette, children, adolescents, morbid obese (BMI above 41Kg/m<sup>2</sup>), bedridden, physically challenged, and subjects above 50 years were excluded from the study.

### 2.5 Ethical Approval and Informed Consent

Ethical approval was sought and obtained from the Research Ethics Committee of the Nnamdi Azikiwe University Teaching hospital (RECNAUTH) Nnewi, Anambra state with reference NAUTH/CS/66/VOL10/2017/010. The participants were informed about the study designs and their written informed consent was obtained before they were recruited.

### 2.6 Data Collection Procedure

Subjects who indicated interest in the study, following discussion at business areas, churches, offices, recreation outlets, and restaurant were given detailed designed questionnaire to fill.

### 2.7 Anthropometric Measurements

The weights of the subjects were evaluated with scale (Gulfex Medical and Scientific, England)). The subjects' heights were recorded in meters using a height scale calibrated in centimeters. As a measure of generalized obesity, each adult participant's BMI was computed by dividing the weight in kilograms, by the square of the height in meters (kg/m<sup>2</sup>). To determine abdominal obesity, measurement of the waist circumference (WC) was taken using a stretch-resistant tape (HTS, China). Blood pressure (BP) systolic and diastolic pressure readings were taken from the participant's left arm using sphygmomanometer

(Omron Medical, United Kingdom). The reading was taken in the morning to the nearest mmHg.

### 2.8 Sample Collection, Storage and Analysis

5mls of blood sample was collected from fasting subjects between 8 and 10am using standard procedure as described by Lewis *et al.*, (2006) [7]. 1 ml of whole blood was dispensed into fluoride oxalate bottle and the plasma separated for glucose analysis while the remaining 4ml of whole blood was dispensed into plain bottle and allowed to clot, retracted and spun at 3000RPM for 10 minutes after which the serum was separated into two aliquots and stored. Plasma glucose was analyzed immediately while serum if not assayed immediately was stored at -20°C not more than 2weeks before analyses. For the cross-sectional study, one point blood sample was collected from each participant both for normal weight and test subjects while in intervention study, three point sample was collected from each test subject: baseline, 6weeks and 12weeks following green tea supplementation. Glucose was assayed colorimetrically using Glucose oxidase method of Trinder, [8]. MDA level was determined by the colorimetric method of Gutteridge and Wilkins, [9]. The serum insulin level was estimated based on solid phase enzyme linked immunosorbent assay (ELISA) method using ACUBIND kit and mindray (MR- 96A) ELISA machine. Insulin Resistance (IR), was assessed by homoeostasis model assessment–insulin-resistance index (HOMA–IR), according to the following formulas: 'fasting insulin value (mU/L) × fasting blood sugar level (mmol/L) / 22.5' [10], values exceeding 2.25 would denote insulin resistance. Quality control was ensured by using pooled control sera from apparently healthy individual and commercially purchased control (Randox (USA) Control level 1.

### 2.9 Statistical Analyses

Statistical analyses were performed using statistical package for social sciences (SPSS) software version 23.0 software. The variables were expressed as mean ± SD. A preliminary comparison of differences between obese Class I, Class II, and non-obese (control), was assessed using Analysis of Variance (ANOVA) while Post Hoc was used for inter-group variability. Paired t-test was used to assess the mean difference between two related variables and level of significant was considered at P<0.05.

**3. RESULTS**

**3.1 Anthropometric Measurement in Obese (Class II and Class I) and Normal Weight Groups (Control)**

The test groups were age matched with the control group, therefore there was no significant difference in the mean age across the groups (P>0.05). (Class II; 38.2±5.26, Class I; 38.95±5.69, normal weight group (control) 36.6 ±5.1). There were significant increases in the mean values of weight, waist circumference, W/H ratio, height, BMI in obese subjects (class II and class I obese) when compared with control group (P<0.05), likewise in Class II obese when compared with Class I obese (P<0.05) except height which did not decrease significantly.

In Table 2, the mean levels of fasting plasma glucose, insulin, HOMA-IR, systolic and diastolic blood pressure increased significantly in obese group (class II and class I) when compared with

their control likewise in Class II obese when compared with Class I obese (P<0.05) while in case of MDA, significant increase was found only in Class II obese subjects when compared with the normal weight subjects.

In Table 3, significant weight loss was observed only after 12 weeks of green tea supplementation when compared with baseline and also at 12weeks when compared with 6weeks of supplementation(P<0.05) unlike waist circumference which reduced significantly after 6 and 12wks intervention when compared with baseline (P <0.05). Furthermore at 12weeks supplementation, there were significant decreases in systolic blood pressure, fasting plasma glucose, fasting blood insulin, MDA, homeostatic model assessment - Insulin resistance (HOMA-IR) (P <0.05) when compared with their baseline values, however FPG, HOMA-IR, MDA and systolic pressure did not significantly decrease after the first 6weeks of intervention (P>0.05).

**Table 1. Anthropometric measurement in Obese (Class II and Class I) and non-obese groups (control) MEAN ± SD**

Parameter	Class II obesity	Class I obesity	Control (normal weight)	F value	P value	Post HOC		
						B/C	A/C	A/B
AGE (yrs.)	38.2±5.26	38.95±5.69	36.6 ±5.1	2.6	0.075	0.087	0.435	1.000
HEIGHT (m)	1.7 ± 0.07	1.72 ±.071	1.75 ±.07	7.6	.001*	.045*	.000*	0.624
WEIGHT(kg)	109±8.35	97 ±8.45	69.7 ±4.6	423.6	.000*	.000*	.001*	.000*
BMI (kg/m <sup>2</sup> )	38.2±1.06	32.9 ±1.06	22.7 ±1.1	2542	.000*	.000*	.000*	.000*
WAIST(cm)	114± 7.28	106 ±5.1	85 ± 9.0	206	.000*	.000*	.000*	.000*
HIP (cm)	111.9±9.7	111.8 ±4.6	96 ± 9.7	59.5	0.000*	0.000*	0.000*	1.000
W/H RATIO	1.03±.059	0.95 ±.05	0.89 ±.07	59.5	0.000*	0.000*	0.000*	0.000*

KEY : A represents class ii obesity, B represents class i obesity, C represents control. BMI = Body mass index, Key \* = Results compared are significantly different at P-value < 0.05 (P < 0.05)

**Table 2. Mean fasting plasma glucose, fasting blood insulin, HOMA-IR and blood pressure in obese subject (Class II and Class I) and non-obese groups (control) MEAN ± SD**

Parameter	Class II Obesity	Class I Obesity	Normal Weight	Fvalue	Pvalue	Post HOC		
						B vs C	A vs C	B vs C
SBP (mm/Hg)	136.9±8.0	130.9±14	123.7±7.8	20.5	.000*	0.012*	0.000*	0.026*
FPG (mmol/L)	6.04 ±.77	5.59 ±.88	5.12 ±.74	16	0.000*	0.035*	0.000*	0.012*
INSULIN (µIU/ml)	7.7 ±2.6	6.3 ±2.2	4.7 ±1.4	26	0.001*	0.019*	0.000*	0.014*
HOMA-IR	2.1 ±.75	1.6 ±.61	1.05 ±.32	39.5	0.000*	0.001*	0.000*	0.000*
DBP(mm/Hg)	95.2±7.1	88 ±9.8	82.9 ±8.9	23	0.000*	0.001*	.000*	0.012*
MDA(nmol/ml)	3.94±1.27	3.72 ±.91	3.30 ±.87	5.1	.007*	0.122	.008*	0.985

Key \* = Results compared are significantly different at P-value < 0.05 (P < 0.05). KEY : A represents class ii obesity, B represents class i obesity, C represents control. FPG= Fasting Plasma Glucose, HOMA-IR Homeostatic Model Assessment-Insulin Resistance, MDA malonaldehyde, SBP, systolic Blood Pressure, DBP Diastolic Blood Pressure

**Table 3. Mean level of blood pressure, fasting blood glucose, fasting blood insulin, MDA and HOMA-IR at different stages of green tea supplementation**

Parameters N= 20	Baseline (A)	6weeks (B)	12week (C)	Post hoc		
				A vs B	A vs C	B vs C
MDA (nmol/L)	3.95±.66	3.83±.97	3.31±.88	0.693	0.018*	0.071
Waist circu. (cm)	112.6±8.9	112.2±9.2	111.9±9.5	0.016*	0.006*	0.297
SBP (mm/Hg)	133.6±8.8	133.5±8.9	132.9±8.6	0.614	0.031*	0.017*
DBP (mm/Hg)	93.5±7.6	93.35±7.84	93.1±7.85	0.614	0.104	0.204
FPG (mmol/L)	5.6±.83	5.4±.77	5.3±.75	0.089	0.003*	0.031*
FBI ( uIU/L)	7.9±1.2	6.0±1.6	5.14±.99	0.000*	0.001*	0.065
HOMA-IR	2.0±.74	2.2±.61	1.6±.47	0.316	0.039*	0.006*

Key \* = Results compared are significantly different at  $P$ -value  $< 0.05$  ( $P < 0.05$ ).

FPG= Fasting Plasma Glucose, HOMA-IR Homeostatic Model Assessment-Insulin Resistance, MDA malondialdehyde, SBP, systolic Blood Pressure, DBP, Diastolic Blood Pressure

#### 4. DISCUSSION

Obesity is becoming one of the most prevalent health concerns among all populations and age groups worldwide. It results to a significant increase in mortality and morbidity related to coronary heart diseases, diabetes type 2, metabolic syndrome, stroke, oxidative stress and cancers [11].

This present study shows that obesity significantly increases fasting plasma glucose, insulin, MDA, blood pressure and it causes insulin resistance ( $P < 0.05$ ). In comparing class II with Class I obese, there were significant increases in FPG, Insulin, HOMA-IR, systolic and diastolic blood pressure ( $P < 0.05$ ) except MDA which did not differ significantly ( $P > 0.05$ ). This findings are in line with those of Gurung et al. [12] which also showed significant increased levels of fasting serum insulin and insulin resistance in obese group when compared with their controls. This indicates that class II obese groups are at the highest risk of developing atherosclerosis, hypertension and diabetes mellitus. This increase might be as a result of the link between obesity and impaired serum glycemic levels resulting from different cellular mechanisms including alterations of insulin signaling, changes in glucose transport, pancreatic  $\beta$  cell dysfunction, as well as enhanced oxidative stress (OS) and inflammation [13]. Obese individuals have demonstrated markers indicative of oxidative stress, including elevated measures of reactive oxygen species (ROS) [14] and diminished antioxidant defense, which is associated with lower antioxidant enzymes as a result of increased free fatty acid which inhibits NADPH oxidase causing dysregulation of cytokines consequently leading to insulin resistance. Oxidative stress is

associated with systemic inflammation, endothelial cell proliferation and apoptosis, and increased vasoconstriction, and thus a noteworthy contributing factor to endothelial dysfunction [15].

In this study, significant weight loss was observed only after 12 weeks of green tea supplementation when compared with baseline value and also at 12weeks when compared with 6weeks of supplementation ( $P < 0.05$ ), however waist circumference reduced significantly after 6 and 12wks of intervention when compared with the baseline value ( $P < 0.05$ ). This is in line with an intervention study by Suzuk et al. [16] which revealed that subjects with high catechin intake had lower body weights, BMI, abdominal circumference and total abdominal fat area, after 12 weeks than those of the placebo group. In contrast, supplementation with 300 mg/d of EGCG for 12 weeks according to Mielgo-Ayuso et al. [17] did not improve weight-loss. However, few others demonstrated that green tea has no effect on FPG [18]; there were no glucose or insulin-lowering effects after consumption of 300 mL of green tea or water [19]. Weight reduction by green tea observed in this study may be due to reduced rate of digestion and an increase in energy expenditure and fat oxidation through  $\beta$ -adrenoceptor activated thermogenesis of brown adipose tissue [20] and also due to inhibition of catechol-O-methyl transferase (COMT) enzyme by epigallocatechingallate (EGCG) of the green tea [21]. Furthermore at 12weeks supplementation in this work, there were significant decreases in systolic blood pressure, fasting plasma glucose, fasting blood insulin, MDA, homeostatic model assessment - Insulin resistance (HOMA-IR) ( $P < 0.05$ ) when compared with their baseline values, however FPG, HOMA-IR, MDA and systolic blood

pressure did not decrease significantly after the first 6 weeks of intervention. ( $P > 0.05$ ) Furthermore, the diastolic blood pressure did not reduce significantly throughout the 12 weeks of supplementation ( $P > 0.05$ ). This finding is in line with the works done by other researchers [22] [23] and also with the work of Liu et al. [24] which also showed that green tea extract caused a significant decrease in homeostasis model assessment of insulin resistance index after 16 weeks of consumption. The anti-hyperglycemic effect of green tea as seen in this study might be as a result of the increase in insulin-stimulated glucose uptake, inhibition of the intestinal GLUT system and decrease in expression of genes that control gluconeogenesis. Mozaffari-Khosravi et al. [25] observed significant decrease in systolic and diastolic blood pressures on individuals who consumed three glasses of green tea daily for 4 weeks, however, in the present study green tea did not reduce the blood pressures at 6 weeks of supplementation. This may be as a result of GT dosage, rate of consumption or the brand of tea used. The decreased blood pressure observed in this study from green tea consumption is probably because it regulates vascular homeostasis by influencing the production of angiotensin II, prostaglandins, endothelin-1 as well as vasodilating substances such as prostacyclin.

## 5. CONCLUSIONS

Obesity is the major cause of diabetes and insulin resistance. Green tea could be beneficial to diabetic patients and obese subjects. Green tea compounds- phytochemicals could be beneficial as one of the components of their diet.

## CONSENT

All authors declare that 'written informed consent was obtained from the subjects and other approved parties for publication of this paper.

## ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee (The Ethical Review Committee of the Nnamdi Azikiwe University Teaching Hospital, Nnewi, Nigeria) and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Adedayo OS, Ebenezer OO. Prevalence of obesity among adolescents in Ile-Ife, Osun state, Nigeria using body mass index and waist hip ratio: A comparative study. 2013; 54(3):153-156.
2. Akiibinu MO, Soile BO, Ajibola M, Amzat U, Olatunji TK. Plasma levels of CA125, CEA, AFP and cortisol in obesity. *Journal of Steroids & Hormonal Science*. 2015; 4:2.
3. Rossner S. Obesity: The disease of the twenty-first century. *International Journal of Obesity Related Metabolic Disorder*. 2002; 26(Suppl 4):S2-4.
4. Mann J, Truswell S. *Essentials of human nutrition*. Oxford Univ. Press. 2012; 69:316.
5. Wu AH, Spicer D, Stanczyk F.Z, Tseng C.C, Yang CS, Pike MC. Effect of 2-month controlled green tea intervention on lipoprotein cholesterol, glucose, and hormone levels in healthy postmenopausal women. *Cancer Prev Respondent*. 2012; 5:393-402.
6. Kelly Da, Nelson R. Characterization and quantification of phytochemicals by gas chromatography. *Journal of Brazil. Chemistry. Society*. 2014;25.
7. Lewis SM, Bain BJ, Bates I, Dacie JV. *Dacie and lewis practical haematology*. Philadelphia Churchill Living Stone; 2006.
8. Trinder P. Determination of blood glucose using an oxidase- peroxidase system with a non carcinogenic chromogen. *Journal of Clinical Pathology*. 1969;22(1):158-161.
9. Gutteridge JM, Wilkins S. Copper dependant hydroxyl radical damage to ascorbic acid; formation of thiobarbituric acid reactive products. *FEBS Letts*. 1982; 137:327-330.
10. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC; Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 1985;28:412-419.
11. Eckel RH, York DA, Rössner S, Hubbard V, Caterson I, St Jeor ST. American Heart Association. Prevention conference VII

- obesity, a worldwide epidemic related to heart disease and stroke: Executive summary. *Circulation*. 2004;110:2968–2975.
12. Gurung A, Meera S, Ebenezer W. Study of insulin resistance and lipid profile in obese adults. *International Journal of Science Innovations and Discoverie*. 2013;3(3): 362-366.
  13. Fernández-Real JM, Broch M, Vendrell J, Ricart W. Insulin resistance, inflammation, and serum fatty acid composition. *Diabetes Care*. 2003;26(5):1362–1368.
  14. Keane JF, Larson MG, Vasani RS, Wilson PW, Lipinska I, Corey D. Obesity and systemic oxidative stress clinical correlates of oxidative stress in the framingham study. *Arteriosclerosis Thrombosis Vascular Biology*. 2003;23:434–9.
  15. Huang C, Matthew J, McAllister, Aaron L, Slusher, Heather E. Webb. Obesity-related oxidative stress: The impact of physical activity and diet manipulation. *Sports Medicine Open*. 2015;1:32.
  16. Suzuki Y, Miyoshi N, Isemura M. Health-promoting effects of green tea. *Proc Jpn Acad Ser B Phys Biol Sci*. 2012;88(3):88–101.
  17. Mielgo-Ayuso J, Barrenechea L, Alcorta P, Larrarte E, Margareto J, Labayan I. Effects of dietary supplementation with epigallocatechin-3-gallate on weight loss, energy homeostasis, cardiometabolic risk factors and liver function in obese women: Randomised, double-blind, placebo-controlled clinical trial. *British Journal and Nutrition*. 2014;111(7):1263–1271.
  18. Mozaffari-Khosravi H, Ahadi Z, Barzegar K. The effect of green tea and sour tea on blood pressure of patients with type 2 diabetes: A randomized clinical trial. *Journal of Diet Supplement* 2013; 10(2):105–115.
  19. Josic J, Olsson AT, Wickeberg J, Lindstedt S, Hlebowicz. Does green tea affect postprandial glucose, insulin and satiety in healthy subjects: A randomized controlled trial? *Nutrition Journal*. 2010;9:63.
  20. Hasani-Ranjbar S, Zahra JH, Mohammad AI. A systematic review of anti-obesity medicinal plants - an update. *Journal of Diabetes and Metabolic Disorder*. 2013; 12:28.
  21. Wang S, Moustaid-Moussa N, Chen L, Mo H, Shastri A, Su R. Novel insights of dietary polyphenols and obesity. *Journal of Nutrition and Biochemistry*. 2014;25(1):1–18.
  22. Peng X. Effect of green tea consumption on blood pressure: A meta-analysis of 13 randomized controlled trials. *Science Report* 1. 2014;4:6251.
  23. Szulińska M, Marta Stępień, Matylda Kręgielska-Narożna, Joanna Suliburska. Effects of green tea supplementation on inflammation markers, antioxidant status and blood pressure in NaCl-induced hypertensive rat model. *Journal of Food & Nutrition Research*. 2017;61(1).
  24. Liu CY, Huang CJ, Huang LH, Chen IJ, Chiu JP, Hsu CH. Effects of green tea extract on insulin resistance and glucagon-like peptide 1 in patients with type 2 diabetes and lipid abnormalities: A randomized, double-blinded, and placebo-controlled trial. *PLoS One*. 2014; 9(3):e91163.
  25. Bhardwaj P, Khanna D. Green tea catechins: Defensive role in cardiovascular disorders. *Chinese Journal of Natural Medicine*. 2013;11:345–353.

© 2018 Joy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sciencedomain.org/review-history/24412>