



## **Evaluation of Growth Performance of Some Rice Varieties in Relation to Their Economic Yield**

**Towfia Junnat<sup>1\*</sup>, Md. Moinul Haque<sup>1</sup>, Md. Elias Hossain<sup>1</sup>, Md. Tofail Hosain<sup>1</sup>,  
Rezowana Nizam<sup>1</sup> and Md. Meftaul Islam<sup>2</sup>**

<sup>1</sup>Department of Agricultural Botany, Sher-e-Bangla Agricultural University, Dhaka, Bangladesh.

<sup>2</sup>Department of Agricultural Chemistry, Sher-e-Bangla Agricultural University, Dhaka, Bangladesh.

### **Authors' contributions**

*This work was carried out in collaboration between all authors. Author TJ designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Author MMH supervised the research work and author MEH managed the analyses of the study. Authors MTH, RN and MMI managed the literature searches. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/ARJA/2017/37394

#### Editor(s):

(1) M. Yuvaraj, Assistant Professor, Department of Soil Science and Agricultural Chemistry, Adhiparasakthi Agricultural College, Tamil Nadu, India

#### Reviewers:

(1) Mahmoud Hozayn Mahmoud, National Research Centre, Egypt.

(2) Eriogun Henry, Federal University of Technology, Nigeria.

(3) Kürşat Çavuşoğlu, Süleyman Demirel University, Turkey.

(4) Nebi Bilir, Süleyman Demirel University, Turkey.

Complete Peer review History: <http://www.sciencedomain.org/review-history/22283>

**Original Research Article**

**Received 13<sup>th</sup> October 2017**  
**Accepted 15<sup>th</sup> November 2017**  
**Published 14<sup>th</sup> December 2017**

### **ABSTRACT**

The experiment was conducted during the period of March to August (*Aus* season), 2013 at the experimental farm of Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, and Dhaka to evaluate the growth performance of some rice varieties in relation to their economic yield. The performance of BRRI hybrid dhan1, BRRI hybrid dhan2, ACI hybrid2, Jagoron and Panna1 were compared with check variety, BRRI dhan48. The experiment was laid out with Randomized Complete Block Design (RCBD) with three replications. Data were recorded on different growth parameters, grain yield, straw yield, biological yield and harvest index. The data were recorded at 30, 45, 60 and 75 days after transplanting (DAT) as well as at final harvest and obtained data were statistically analyzed for comparisons. In term of vegetative growth parameters, plant height, tiller hill<sup>-1</sup>, leaf area index, stem dry matter hill<sup>-1</sup>, leaves dry matter hill<sup>-1</sup>

\*Corresponding author: E-mail: towfijannat@yahoo.com;

and total dry matter  $\text{hill}^{-1}$  at different DAT, the ACI hybrid2 was superior over the check variety, BRR1 dhan48 as well as BRR1 hybrid dhan1 and BRR1 hybrid dhan1. Highest and statistically significant grain yield ( $3.05 \text{ t ha}^{-1}$ ) straw yield ( $4.20 \text{ t ha}^{-1}$ ) biological yield ( $7.25 \text{ t ha}^{-1}$ ) and harvest index (42.01%) was observed in ACI hybrid2 which was statistically identical with hybrid rice Jagoron and Panna. All test hybrid varieties were superior in vegetative and yield performance over inbreed BRR1 dhan48 during *Aus* season.

**Keywords:** Rice; hybrid variety; *Aus* season; economic yield.

## 1. INTRODUCTION

Rice (*Oryza sativa* L.) is an important food around the world and the staple food approximately more than two billion people in Asia [1]. The population of Bangladesh is increasing at an alarming rate and the cultivable land is decreasing due to urbanization and industrialization resulting in more shortage of food. The nation is still adding about 2.3 million every year to its total of 150 million people [2]. Thus, the present population is supposed to swell progressively to 223 million by the year of 2030 which will require additional 48 million tons of food grains instead of current deficit of about 1.2 million tons every year [3]. So, the highest priority has been given to increase its production per unit area [4].

In Bangladesh, the geographical, climatic and edaphic conditions are favorable for year round rice cultivation. But the national average rice yield in Bangladesh is only  $2.57 \text{ t ha}^{-1}$  [5]. The average rice yield of rice in Bangladesh is far below compared to other rice growing countries, like China is ( $8.75 \text{ t ha}^{-1}$ ), Japan ( $8.22 \text{ t ha}^{-1}$ ) and Korea ( $8.04 \text{ t ha}^{-1}$ ) [6]. In Bangladesh, rice dominates over all other crops and covers 75% of the total cropped area of which around 79% is occupied by high yielding rice varieties [7].

Rice yield can be increased in many ways. Horizontal expansion of rice growing area, development of new high yielding and hybrid varieties, introduction of developed varieties, adaption of proper agronomic management practices can help to achieve potential yield per unit area and total production of rice. It has potentiality for food security of poor countries where arable land is scarce populations is expanding and labor is cheap. In our country, Bangladesh Rice Research Institute (BRR1) has started breeding program for the development of super high yielding varieties with large panicles and high yield potentialities.

Rowth and yield of rice are strongly influenced by genotype as well as environmental factors [8,9]. The genetic potentiality of a rice variety is almost fixed, but grain yield can be increased by the manipulation of management practices and by growing in recommended season [10]. Now a day's different hybrid rice varieties are available in Bangladesh which have more yield potential than conventional high yielding varieties [11]. Improvement of rice grain yield is the main target of breeding program to develop rice varieties for diverse ecosystems. In addition, grain yield also related with other characters such as growth duration, and yield components [12].

There are three distinct rice growing seasons in Bangladesh namely *Aus*, *Aman* and *Boro*. Among these three, *Aus* rice covers only 12.27% of the rice growing area with average yield is  $1.73 \text{ t ha}^{-1}$  [5]. Recently various new high yielding rice varieties have been developed by BARI for *Aus* season like BRR1 dhan42, BRR1 dhan48, BRR1 dhan55 etc. but there yield is much lower than hybrid rice varieties, because the hybrids have larger panicles and heavier seeds resulting an average grain yield of  $7.27\%$  [13]. The low yield of rice in the *Aus* season should be improved to meet the current demand of staple food and future food security. Hybrid rice is one of the viable options for increasing the yield over the best modern varieties. Based on above proposition, this research work was designed to evaluate the growth and yield performance of some hybrid rice varieties in *Aus* season.

## 2. MATERIALS AND METHODS

The current research work was conducted in the experimental field of Sher-e-Bangla Agricultural University, Dhaka during *Aus* season (March to August) of 2013. Geographically this location is under the subtropical climate, characterized by three distinct seasons, winter season from November to February and the pre-monsoon period or hot season from March to April and monsoon period from May to October [14].

The soil of the experimental area belongs to "The Modhupur Tract", AEZ-28 [15]. Top soil was silty clay in texture, olive-gray with common fine to medium distinct dark yellowish brown mottles. Soil pH was 5.6 and has organic carbon 0.45%. The experimental area was flat having available irrigation and drainage system. The experiment comprised of six rice varieties which were used as treatment for the experiment viz., V1: BRR1 dhan48, V2: BRR1 hybrid dhan1, V3: BRR1 hybrid dhan2, V4: ACI hybrid2, V5: Jagoron and V6: Panna. The experiment was laid out in Randomized Complete Block Design (RCBD) with three replications. All the rice varieties used in this study were hybrid except BRR1 dhan48. Among the varieties, BRR1 dhan48, BRR1 hybrid dhan1 and BRR1 hybrid dhan2 were released by BRR1, Joydebpur, Dhaka. The introduced Chinese varieties, ACI hybrid2, Jagoron and Panna1 were imported and marketed by ACI Ltd., Bangladesh. Seeds of these varieties were collected from respective organizations.

Healthy seeds were selected by specific gravity method and then immersed in water bucket for 24 hours and then they were kept tightly in gunny bags. The seeds started sprouting after about 48 hours and were sown after 72 hours. Seeds of the selected cultivars were sown separately in well prepared 2m x 1m sized wet seedbed for each cultivar on march 10, 2013. Recommended doses of fertilizers and manures were applied during seedbed preparation.

The selected plot was ploughed with a power tiller for sun dry in April 6, 2013 and left for a week. After one week the land was irrigated, harrowed, ploughed and cross-ploughed several times followed by laddering to obtain desirable tilth of soil for transplanting seedlings. After removal of weeds and stubbles the plot was partitioned into unit plots in accordance with the experimental design. The fertilizers N, P, K, S, Zn and B were applied in the form of urea, Triple Super Phosphate (TSP), Muriate of Potash (MOP), gypsum, zinc sulphate and borax, respectively. The entire amount of TSP, MOP, gypsum, zinc sulphate and borax were applied during the final preparation of plot land. Recommended dose of Furadan 57 EC and decomposed cowdung @ 10 t ha<sup>-1</sup> were applied during land preparation. Urea was applied in two equal installments at tillering and before panicle initiation. The rice seedlings were uprooted on April 09, 2013. The nursery bed

was irrigated a day before uprooting of the seedlings to minimize mechanical injury of roots. Seedlings were transplanted in main plots having 30 cm x 25 cm spacing between line to line and plant to plant in April 10, 2013. Two healthy seedlings were transplanted in each hill.

After establishment of seedlings, various intercultural operations were accomplished for better growth and development of the rice seedlings. Gap filling was executed for all of the plots at 10 days after transplanting (DAT) by planting same aged seedlings. In the early stage of growth flood irrigation was provided to maintain a constant level of standing water up to 6 cm to enhance tillering and 10-12 cm in the later stage to depress late tillering. The newly emerged weeds were uprooted carefully at tillering stage and before panicle initiation. The field was finally dried out 15 days before harvesting.

The rice was harvested manually from each plot depending upon the maturity. The harvested crop of each pot was bundled separately, properly tagged and brought to threshing floor. The threshing and cleaning of rice seed were conducted with great care. Plot wise fresh weight of grain and straw were recorded. Finally the weight of grains was adjusted to a moisture content of 14%. The straw was sun dried and the yields of grain and straw plot<sup>-1</sup> were recorded.

## 2.1 Data Collection

Field data for plant height, tiller hill<sup>-1</sup>, leaf area index, stem dry matter, tiller hill<sup>-1</sup>, leaves dry matter hill<sup>-1</sup>, stem dry matter hill<sup>-1</sup>, total dry matter hill<sup>-1</sup> were collected at 30, 45, 60 and 75 days after transplanting (DAT). The height of plant at harvest was also recorded. The height was measured from the ground level to the tip of the sample plants. The number of tillers hill<sup>-1</sup> was recorded from each plot by counting total tillers of five random hills and average value was recorded. The leaf area index was measured manually from five randomly selected plants of inner rows of each plot. The final data calculated multiplying by a correction factor 0.75 [12]. For stem and leaves dry matter hill<sup>-1</sup>, three hills was randomly collected from inner rows of each plot. Collected hills were oven dried at 70°C for 72 hours then transferred into desiccator to cool down at room temperature. The desiccated stems and leaves were separated, weighted with electric balance as well

as mean values hill<sup>-1</sup> recorded. Total dry matter hill<sup>-1</sup> was recorded by adding values of stem dry matter hill<sup>-1</sup> and leaves dry matters hill<sup>-1</sup>. Rice grains obtained from harvested area of plot were sun-dried and adjusted to 14% moisture content using a digital moisture meter. Then dried grains of each harvested area were measured carefully with electric balance and measured values were converted to t ha<sup>-1</sup>. Rice straw of obtained from each unit plot were sun-dried and weighed carefully. The sub-samples of the straw of each plot were oven dried, weighted and finally converted to t ha<sup>-1</sup>. Grain yield and straw yield together were regarded as biological yield. Harvest index was calculated from the grain and straw yield of rice for each plot and expressed in percentage.

## 2.2 Statistical Analysis

The data obtained were statistically analyzed using MSTAT software to observe the significant difference among the different rice varieties. The significance of the difference among the treatment means was estimated by the Least Significant Difference Test (LSD) test at 5% level of probability [16].

## 3. RESULTS AND DISCUSSION

### 3.1 Plant Height

A statistically significant variation was observed in plant height of the hybrid rice varieties at different DAT (Table 1). At 30, 45, 60, 75 DAT and harvest, the tallest plant (35.87, 50.40, 76.50, 95.77 and 139.97 cm, respectively) was recorded from ACI hybrid2 which was statistically similar (35.00, 49.03, 76.20, 92.97 and 110.40 cm, respectively) with the height of Jagoron, whereas the shortest plant (29.33,

41.73, 69.03, 82.57 and 102.93 cm, respectively) was found in BRR1 dhan48. Plant height varies differentially on the basis of their varietal characters and improved varieties are the first and foremost requirement for initiation and accelerated production program. Mondal et al., reported that significant variation in plant height exist among test rice varieties in Aus season [17]. Growth of rice is strongly influenced by genotype as well as environmental factors [8].

### 3.2 Tillers Hill<sup>-1</sup>

Statistically significant variation was recorded in terms of tillers hill<sup>-1</sup> due to different rice variety at 30, 45, 60 and 75 DAT (Fig. 1). At 30, 45, 60 and 75 DAT, the maximum tillers hill<sup>-1</sup> (5.67, 8.80, 15.70 and 16.80, respectively) was recorded from ACI hybrid2 which was similar with Jagoron. On the other hand, the minimum tillers hill<sup>-1</sup> (4.07, 6.27, 9.57 and 12.03, at 30, 45, 60 and 75 DAT respectively) was observed from BRR1 dhan48 which was statistically similar with BRR1 hybrid dhan2 and BRR1 hybrid dhan1. Significant variation in tiller hill<sup>-1</sup> was also reported by Mondal et al. [17].

### 3.3 Leaf Area Index

Leaf area index of different rice varieties varied significantly at 30, 45, 60 and 75 DAT. The leaf area index of ACI hybrid2 (0.25, 1.18, 3.64 and 5.91 at 30, 45, 60 and 75 DAT respectively) was significantly higher and statistically similar with Jagoron at same DAT. The lowest leaf area index (0.20, 0.83, 2.64 and 5.13 at 30, 45, 60 and 75 DAT respectively) was found in BRR1 dhan48 which was statistically similar with BRR1 hybrid dhan2 (Table 2).

**Table 1. Plant height at different DAT and harvest of some selected rice varieties**

Rice variety	Plant height (cm)				
	30 DAT	45 DAT	60 DAT	75 DAT	Harvest
BRR1 dhan48	29.33 b	41.73 b	69.03 c	82.57 c	102.93 b
BRR1 hybrid dhan1	30.30 b	43.30 b	70.20 bc	88.53 abc	106.77 b
BRR1 hybrid dhan2	29.43 b	42.57 b	69.07 c	86.03 bc	106.30 b
ACI hybrid2	35.87 a	50.40 a	76.50 a	95.77 a	139.97 a
Jagoron	35.00 a	49.03 a	76.20 a	92.97 ab	110.40 b
Panna1	31.00 b	45.47 ab	73.30 ab	89.90 abc	105.50 b
LSD <sub>(0.05)</sub>	3.675	5.426	3.733	8.066	9.764
Level of significance	0.01	0.05	0.01	0.05	0.01
CV(%)	6.35	6.57	4.84	4.97	4.79

*In a column means having similar letter(s) are statistically similar and those having dissimilar letter(s) differ significantly at 0.05 level of probability*




Fig. 1. Effect of rice varieties in tillers production of tiller hill<sup>-1</sup> at different DAT

### 3.4 Stem and Leaves Dry Matter hill<sup>-1</sup>

Significant variation was observed in terms of stem and leaves dry matter hill<sup>-1</sup> at different DAT of the rice varieties studied (Table 3). At 30, 45, 60 and 75 DAT, the highest stem dry matter hill<sup>-1</sup> (2.63, 3.79, 6.16 and 5.85g, respectively) was recorded from ACI hybrid2 which was statistically similar with Jagoron. The lowest stem dry matter hill<sup>-1</sup> (1.98, 2.98, 4.53 and 4.37 g at 30, 45, 60 and 75 DAT respectively) was observed from BRRi dhan48 which was statistically similar with BRRi hybrid dhan2 and BRRi hybrid dhan1. Data revealed that ACI hybrid2 accumulated maximum leaves dry matter hill<sup>-1</sup> (1.58, 2.83, 5.39 and 5.04 g at 30, 45, 60 and 75 DAT, respectively) and statistically similar with Jagoron at same days after transplanting. On the other hand, BRRi dhan48 and BRRi hybrid dhan2 accumulated the lowest and statistically insignificant amount of leaves dry matter hill<sup>-1</sup> at 30, 45, 60 and 75 DAT (Table 3).

### 3.5 Total Dry Matter Hill<sup>-1</sup>

Rice varieties showed significant variation in total dry matter hill<sup>-1</sup> under the present trial. At 30, 45, 60 and 75 DAT, the highest total dry matter hill<sup>-1</sup> (4.21, 6.62, 11.55 and 10.89 g, respectively) was observed in ACI hybrid2 which was statistically similar with Jagoron but varied with other varieties. The lowest total dry matter hill<sup>-1</sup> (3.21, 5.14, 8.653 and 8.43 g at 30, 45, 60 and 75 DAT respectively) was found in BRRi dhan48 which was statistically similar with BRRi hybrid dhan2 at same data recording date (Table 4). Amin et al. reported that traditional varieties accumulated higher amount of vegetative dry matter than the modern variety [18].

### 3.6 Grain Yield

Rice varieties of the current study showed significant variation in grain yield (Table 5). The highest grain yield (3.05 t ha<sup>-1</sup>) was obtained

Table 2. Leaf area index at different DAT of some selected rice varieties

Rice variety	Leaf area index			
	30 DAT	45 DAT	60 DAT	75 DAT
BRRi dhan48	0.202 c	0.836 c	2.64 c	5.13 b
BRRi hybrid dhan1	0.222 b	0.955 bc	3.11 abc	5.12 b
BRRi hybrid dhan2	0.204 c	0.922 bc	2.96 bc	5.03 b
ACI hybrid2	0.254 a	1.188 a	3.64 a	5.91 a
Jagoron	0.243 ab	1.067 ab	3.44 ab	5.68 ab
Panna1	0.232 b	1.024 ab	3.15 abc	5.60 ab
LSD <sub>(0.05)</sub>	0.058	0.163	0.592	0.721
Level of significance	0.05	0.01	0.05	0.05
CV(%)	12.27	8.88	10.29	7.32

In a column means having similar letter(s) are statistically similar and those having dissimilar letter(s) differ significantly at 0.05 level of probability.

**Table 3. Stem dry matter hill<sup>-1</sup> at different DAT and at harvest of some selected hybrid rice varieties**

Rice varieties	Stem dry matter hill <sup>-1</sup> (g)				Leaves dry matter hill <sup>-1</sup> (g)			
	30 DAT	45 DAT	60 DAT	75 DAT	30 DAT	45 DAT	60 DAT	75 DAT
BRR1 dhan48	1.98 c	2.98 c	4.53 c	4.37 c	1.23 c	2.17 c	4.12 b	4.06 c
BRR1 hybrid dhan1	2.12 bc	3.33 abc	5.51 ab	5.40 ab	1.45 ab	2.59 ab	4.19 b	4.07 c
BRR1 hybrid dhan2	2.08 c	3.28 bc	5.07 bc	5.02 bc	1.39 abc	2.49 b	4.27 b	4.24 bc
ACI hybrid2	2.63 a	3.79 a	6.16 a	5.85 a	1.58 a	2.83 a	5.39 a	5.04 a
Jagoron	2.46 ab	3.47 ab	5.82 a	5.58 ab	1.56 ab	2.70 ab	4.75 ab	4.63 ab
Panna1	2.15 bc	3.36 abc	5.60 ab	5.22 b	1.37 bc	2.43 bc	4.61 ab	4.38 bc
LSD <sub>(0.05)</sub>	0.257	0.438	0.630	0.521	0.191	0.304	0.787	0.386
Level of significance	0.01	0.05	0.01	0.01	0.05	0.01	0.05	0.01
CV(%)	6.32	7.16	6.36	6.67	7.40	6.63	9.52	6.20

*In a column means having similar letter(s) are statistically similar and those having dissimilar letter(s) differ significantly at 0.05 level of probability*

from ACI hybrid2 which was statistically similar with Jagoron (2.97 t ha<sup>-1</sup>) and Panna1 (2.82 t ha<sup>-1</sup>). The lowest grain yield (2.51 t ha<sup>-1</sup>) was observed in BRR1 dhan48 which was statistically identical with BARI hybrid dhan1 and BARI hybrid dhan2. Grain yield is a polygenic character and greatly influenced by environmental factors. Varieties play an important role in producing high yield of hybrid rice. International Rice Research Institute (IRRI) started super rice breeding program to give up to 30% more rice yield (13 to 15 t ha<sup>-1</sup>) than the current modern high yielding plant types [19]. In 2004, Akbar reported that available hybrid rice varieties of Bangladesh have more yield potential than conventional high yielding varieties [11]. Wang et al. reported that compared with conventional cultivars, the hybrids had larger panicles, heavier seeds, resulting in an average yield increase of 7.27%

[13]. Varietal influence of rice on grain yield was also reported [20].

### 3.7 Straw Yield, Biological Yield and Harvest Index

The test varieties of rice varied significantly in straw yield, biological yield and harvest index (Table 5). The highest straw yield (4.20 t ha<sup>-1</sup>) was observed in ACI hybrid2 which was statistically similar with BRR1 hybrid dhan1, BRR1 hybrid dhan2, Jagoron and Panna1 but varied significantly with BRR1 dhan48. The highest biological yield (7.25 t ha<sup>-1</sup>) was found from ACI hybrid2 which was statistically similar and with Jagoron (7.12 t ha<sup>-1</sup>) and Panna1 (6.84 t ha<sup>-1</sup>). The lowest biological yield (6.38 t ha<sup>-1</sup>) was found from BRR1 dhan48 which was statistically identical with BRR1 hybrid dhan1. The highest harvest index (42.07%) was recorded from

**Table 4. Total dry matter hill<sup>-1</sup> at different DAT and at harvest of some selected hybrid rice varieties**

Rice variety	Total dry matter hill <sup>-1</sup> (g)			
	30 DAT	45 DAT	60 DAT	75 DAT
BRR1 dhan48	3.21 c	5.14 c	8.65 c	8.43 c
BRR1 hybrid dhan1	3.57 bc	5.92 ab	9.61 bc	9.47 b
BRR1 hybrid dhan2	3.47 c	5.77 bc	9.34 bc	9.26 bc
ACI hybrid2	4.21 a	6.62 a	11.55 a	10.89 a
Jagoron	3.92 ab	6.17 ab	10.56 ab	10.21 ab
Panna1	3.52 c	5.80 bc	10.20 b	9.60 bc
LSD <sub>(0.05)</sub>	0.382	0.700	1.258	0.840
Level of significance	0.01	0.01	0.01	0.01
CV(%)	5.78	6.53	6.92	5.97

*In a column means having similar letter(s) are statistically similar and those having dissimilar letter(s) differ significantly at 0.05 level of probability*

**Table 5. Effect of rice varieties on grain yield, straw yield, biological yield and harvest index**

Rice variety	Grain yield (t ha <sup>-1</sup> )	Straw yield (t ha <sup>-1</sup> )	Biological Yield (t ha <sup>-1</sup> )	Harvest index (%)
BRRi dhan48	2.51 b	3.87 b	6.38 b	39.34 b
BRRi hybrid dhan1	2.66 b	4.11 a	6.77 b	39.29b
BRRi hybrid dhan2	2.75 b	4.05 a	6.80 b	40.44ab
ACI hybrid2	3.05 a	4.20 a	7.25 a	42.07 a
Jagoron	2.97 a	4.15 a	7.12 a	41.71 a
Panna1	2.82 a	4.02 a	6.84 a	41.23 a
LSD <sub>(0.05)</sub>	0.281	0.182	0.486	1.478
Level of significance	0.05	0.05	0.05	0.05
CV(%)	4.61	5.12	4.04	5.11

*In a column means having similar letter(s) are statistically similar and those having dissimilar letter(s) differ significantly at 0.05 level of probability*

ACI hybrid2 which was statistically similar with Jagoron and Panna1 whereas the lowest harvest index (39.34%) was obtained from BRRi dhan48 and was statistically similar with BRRi hybrid dhan1 and BRRi hybrid dhan2. Patel [21] reported that mean yield increased with Kranti over IR36 was 10.0% for straw yield. Mondal et al. [17] also found significant variation in straw yield, biological yield and harvest index among the test rice varieties in Aus season.

#### 4. CONCLUSION

ACI hybrid2 produced the tallest plant, the maximum number of tillers hill<sup>-1</sup> the highest leaf area index, the highest stem dry matter hill<sup>-1</sup>, the highest leave dry matter hill<sup>-1</sup> and the highest total dry matter hill<sup>-1</sup>, which was statistically significant over check variety, BRRi dhan48 as well as BRRi hybrid dhan1 and BRRi hybrid dhan2 but statistically similar with hybrid rice Jagoron and Panna1. The highest grain yield (3.05 t ha<sup>-1</sup>) was obtained from ACI hybrid2 which was statistically similar with Jagoron and Panna1. The lowest grain yield (2.51 t ha<sup>-1</sup>) was obtained from BRRi dhan48. The highest straw yield (4.20 t ha<sup>-1</sup>) was observed from ACI hybrid2 whereas the straw yield (3.87 t ha<sup>-1</sup>) was lowest in BRRi dhan48. The highest biological yield (7.25 t ha<sup>-1</sup>) was found in ACI hybrid2, while the lowest biological yield (6.38 t ha<sup>-1</sup>) in BRRi dhan48. The highest harvest index (42.07%) was recorded from ACI hybrid2, whereas the lowest (39.34%) was from BRRi dhan48. In term of grain yield, straw yield, biological yield and harvest index the ACI hybrid2, Jagoron and Panna1 were statistically indifferent. However, vegetative and yield performance of all five test hybrid varieties were superior over the inbred variety BRRi dhan48. Above findings revealed that the ACI hybrid2

was superior compared to other varieties in term of growth parameters and yield of rice followed by hybrid rice variety Jagoron.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Hien NL, Yoshihashi T, Sarhadi WA. Evaluation of aroma in rice (*Oryza sativa* L.) using KOH method, molecular markers and measurement of 2-acetyl-1-pyrroline concentration. Japanese. J. Trop. Agron. 2006;50:190-198.
- Momin SI, Husain M. Technology development and dissemination to augment rice production in Bangladesh. In: 'The Guardian'. 2009;33-35.
- Julfiquar AW, Haque MM, Haque AKGME, Rashid MA. Current status of hybrid rice research and future program in Bangladesh. Proc. Workshop on use and development of hybrid rice in Bangladesh, held at BARC, 18-19, May, 2008.
- Bhuiyan NI. The hybrid rice program for Bangladesh. In: 'Hybrid Rice in Bangladesh: Progress and Future Strategies'. Bangladesh Rice Res. Inst. Publication No. 138. 2004;3-5.
- BBS. Statistical Yearbook of Bangladesh. Bangladesh Bureau of Statistics. Statistics Division, Ministry of Planning, Govt. of Peoples Republic of Bangladesh. Dhaka. Bangladesh. 2010;71.
- FAO (Food and Agriculture Organization). FAO Production Yearbook, Food and Agriculture Organization, Rome, Italy. 2009;56-77.

7. BBS. Yearbook of agricultural statistics of Bangladesh. Bangladesh Bureau of Statistics. Stat. Div., Min. Plan. Govt. of the Peoples Repub. of Bangladesh. 2008;125-127.
8. BRRRI (Bangladesh Rice Research Institute). (2003). Annual Internal Review, held on 19-23 October, 2003. Grain Quality and Nutrition Division. Bangladesh Rice Res. Inst. p. 1-20.
9. Jennings I, Juliano BO, Duff D. Rice grain quality as an emerging priority in National rice breeding program. In: rice grain marketing and quality issues. Los Banos, Laguna, IRRI. 1979;55-64.
10. BRRRI (Bangladesh Rice Research Institute). Annual Report for January 1998- June 1999. Bangladesh Rice Res. Inst. Gazipur, Pub. 135. 1999;9-35, 243.
11. Akbar MK. Response of hybrid and inbred rice varieties to different seedlings ages under system of rice intensification in transplant aman season. M.S. (Ag.) Thesis Dept. Agron. B.A.U, Mymensingh; 2004.
12. Yoshida S. Fundamentals of rice crop science. International Rice Research Institute, Los Banos Manila, Philippines. 1981;269.
13. Wang JL, Xu ZJ, Yi XZ. Effects of seedling quantity and row spacing on the yields and yield components of hybrid and conventional rice in northern China. Chinese J. Rice Sci. 2006;20(6):631-637.
14. Edris KM, Islam ATMT, Chowdhury MS, Haque AKMM. Detailed soil survey of Bangladesh, Dept. Soil Survey, Govt. People's Republic of Bangladesh; 1979.
15. FAO (Food and Agriculture Organization). Production Year Book. Food and Agricultural Organizations of the United Nations Rome, Italy. 1988;42: 190-193.
16. Gomez KA, Gomez AA. Statistical procedure for agricultural research (2<sup>nd</sup> edn.). Int. Rice Res. Inst., A Willey Int. Sci. 1984;28-192.
17. Mondal B, Sarkar MAR, Paul SK, Zaman F. Effect of variety and weeding regime on the yield components and yield of Aus rice. J. Bangladesh Agril. Univ. 2013;11(1): 17-21.
18. Amin MR, Hamid A, Choudhury RU, Raquibullah SM, Asaduzzaman M. Nitrogen fertilizer Effect on tillering, dry matter production and yield of traditional varieties of Rice. Int. J. Sustain. Crop Prod. 2006;1(1):17-20.
19. IRRI (International Rice Research Institute). IRRI Almanac Intl. Rice Res. Inst. Los Bonos, Philippines. 1993;110.
20. Xie W, Wang G, Zhang Q. Potential production simulation and optimal nutrient management of two hybrid rice varieties in Jinhua, Zhejiang Province. J. Zhejiang Univ. Sci. 2007;8(7):486-492
21. Patel JR. Effect of water regime, variety and blue green algae on rice (*Oryza sativa*). Indian J. Agron. 2000;45(1):103-106.

© 2017 Junnat et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<http://sciencedomain.org/review-history/22283>